

FUNCTION THEORY AND HOLOMORPHIC MAPS ON SYMMETRIC PRODUCTS OF PLANAR DOMAINS

DEBRAJ CHAKRABARTI AND SUSHIL GORAI

ABSTRACT. We show that the $\bar{\partial}$ -problem is globally regular on a domain in \mathbb{C}^n , which is the n -fold symmetric product of a smoothly bounded planar domain. Remmert-Stein type theorems are proved for proper holomorphic maps between equidimensional symmetric products and proper holomorphic maps from cartesian products to symmetric products. It is shown that proper holomorphic maps between equidimensional symmetric products of smooth planar domains are smooth up to the boundary.

1. INTRODUCTION

1.1. Symmetric Products. Let $n \geq 2$ be an integer. Given an object X , a basic construction in mathematics is to form its n -fold cartesian product X^n with itself, the set of ordered n -tuples (x_1, x_2, \dots, x_n) of elements $x_i \in X$. One can equally consider the space $X^{(n)}$ of *unordered* n -tuples of elements of X , called the n -fold *symmetric product* of X . If σ is a bijection of the set $I_n = \{1, 2, \dots, n\}$ with itself, the tuples (x_1, x_2, \dots, x_n) and $(x_{\sigma(1)}, x_{\sigma(2)}, \dots, x_{\sigma(n)})$ are to be considered identical in $X^{(n)}$. Unlike in the case of the cartesian product, the smoothness of X is usually not inherited by $X^{(n)}$. For example, if X is a complex manifold, the symmetric product $X^{(n)}$ is in general only a complex space, which locally looks like a complex analytic set, possibly with singularities.

If U is a Riemann surface, then the symmetric product $U^{(n)}$ is a complex manifold of dimension n in a natural way. In particular, if U is a domain in the complex plane, the symmetric product $U^{(n)}$ is in fact biholomorphic to a domain $\Sigma^n U$ in \mathbb{C}^n , which may be constructed in the following way. For $k \in I_n = \{1, 2, \dots, n\}$, denote by π_k the k -th elementary symmetric polynomial in n variables. Let $\pi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be the polynomial map given by $\pi = (\pi_1, \dots, \pi_n)$, referred to as the *symmetrization* map on \mathbb{C}^n . We then define $\Sigma^n U = \pi(U^n)$, where $U^n \subset \mathbb{C}^n$ is the n -fold cartesian product of the domain U . Then $\Sigma^n U$ is biholomorphic to the n -fold symmetric product $U^{(n)}$ (see Section 2.1 below.)

In this paper, we study the domains $\Sigma^n U$ from the point of view of the classical theory of functions and mappings of several complex variables. For any domain $U \subset \mathbb{C}$, $\Sigma^n U$ is a pseudoconvex domain in \mathbb{C}^n . The main interest is therefore in the study of boundary properties of functions and mappings. The boundary $\mathfrak{b}\Sigma^n U$ of $\Sigma^n U$ is a singular hypersurface in \mathbb{C}^n , with a Levi-flat smooth part. Even when U has smooth boundary, the boundary $\mathfrak{b}\Sigma^n U$ is not Lipschitz, a fact which was shown to us recently by Łukasz Kosiński in the case $n = 2$, and from which the general case follows. In spite of having such non-smooth boundary, it turns out that the complex-analytic properties of $\Sigma^n U$ are in many ways similar to those of the cartesian product U^n .

1.2. Main Results. Recall that the $\bar{\partial}$ -problem on a domain consists of solving the equation $\bar{\partial}u = g$, where g is a given (p, q) -form with $q > 0$ such that $\bar{\partial}g = 0$, and u is an unknown $(p, q - 1)$ -form. Since $\Sigma^n U$ is pseudoconvex, the $\bar{\partial}$ -problem always has a solution on it. We also have interior regularity: if the datum g has coefficients which are \mathcal{C}^∞ -smooth, there is a solution u which also has smooth coefficients (see e.g. [28].) The interesting question regarding $\Sigma^n U$ is therefore that of boundary regularity of the $\bar{\partial}$ -problem. Recall that on a domain $\Omega \Subset \mathbb{C}^n$, the $\bar{\partial}$ -problem is said to be *globally regular* if for any $\bar{\partial}$ -closed form $g \in \mathcal{C}_{p,q}^\infty(\bar{\Omega})$ with $q > 0$, there exists a form $u \in \mathcal{C}_{p,q-1}^\infty(\bar{\Omega})$ such that $\bar{\partial}u = g$, where $\mathcal{C}_{p,q}^\infty(\bar{\Omega})$ denotes the

2010 *Mathematics Subject Classification.* 32A07, 32W05, 32H40.

Sushil Gorai was supported in part by an INSPIRE Faculty Fellowship awarded by DST, Government of India.

space of forms of degree (p, q) with coefficients which are smooth up to the boundary. Our first result is the following:

Theorem 1.1. *If $U \Subset \mathbb{C}$ is a domain with C^∞ -smooth boundary in the plane, the $\bar{\partial}$ -problem is globally regular on $\Sigma^n U$.*

For smoothly bounded domains, a classical approach to the study of the regularity of the $\bar{\partial}$ -problem is to reduce it to the study of the $\bar{\partial}$ -Neumann problem for the complex Laplace operator (see [38, 16, 26] for details.) Indeed, from Kohn's weighted theory of the $\bar{\partial}$ -Neumann operator (see [29]) it follows that on a smoothly bounded pseudoconvex domain in \mathbb{C}^n , the $\bar{\partial}$ -problem is globally regular. However, as we show in Proposition 5.3 below, the domain $\Sigma^n U$ is not Lipschitz, even though U may have C^∞ boundary. Consequently many of the basic tools of the theory of the $\bar{\partial}$ -Neumann problem, including the crucial method of "elliptic regularization", do not apply to $\Sigma^n U$. Furthermore, the presence of complex hypersurfaces in the smooth part of $b\Sigma^n U$ means that the $\bar{\partial}$ -Neumann operator on $\Sigma^n U$ is non-compact, and one would even expect it to be highly non-regular (cf. [24].)

For non-smooth pseudoconvex domains, the global regularity of the $\bar{\partial}$ -problem is a subtle matter. It is well-known that on the Hartogs Triangle, the non-Lipschitz domain in \mathbb{C}^2 given by $\{|z_1| < |z_2| < 1\}$, the $\bar{\partial}$ -problem is not globally regular (see [14, 13].) However the geometry of the Hartogs Triangle is very different from that of symmetric products in that the Hartogs Triangle does not admit a basis of Stein neighborhoods. The presence of a Stein neighborhood basis with appropriate geometric properties (" $s - H^n$ -convexity in the terminology of [15]) is crucial in the proof of Theorem 1.1. Note that though the $\bar{\partial}$ -Neumann operator on $\Sigma^n U$ is non-compact, the $\bar{\partial}$ -problem is globally regular. The situation on $\Sigma^n U$ is therefore analogous to that on cartesian products (see [12, 33].)

Next, we study proper holomorphic mappings of symmetric products. It is natural to study mappings between symmetric and cartesian products. When $U = V = \mathbb{D}$, the unit disc in the plane, this was done by Edigarian and Zwonek in [22, 23], using the classical method of Remmert and Stein (see [37, 35].) Generalizing this, we have the following:

Theorem 1.2. *Let U_1, \dots, U_n be bounded domains in the complex plane and let $G = U_1 \times \dots \times U_n$ be their cartesian product. Let $f : G \rightarrow \Sigma^n V$ be a proper holomorphic map. Then there are proper holomorphic maps $g_j : U_j \rightarrow V$ such that*

$$f = \pi \circ (g_1 \times \dots \times g_n),$$

where by definition, for $z \in G$, we have $(g_1 \times \dots \times g_n)(z) = (g_1(z_1), \dots, g_n(z_n))$.

This can be compared with the fact that a proper holomorphic mapping between equidimensional cartesian product domains splits as product of mappings in the factors (see [35, page 77].)

Theorem 1.2 leads to the classification of proper holomorphic maps between equidimensional symmetric products: for any mapping $\psi : U \rightarrow V$ of planar domains, let $\Sigma^n \psi : \Sigma^n U \rightarrow \Sigma^n V$ be the unique map such that

$$(\Sigma^n \psi)(\pi(z_1, \dots, z_n)) = \pi(\psi(z_1), \dots, \psi(z_n)). \quad (1.1)$$

(see Section 2.3 below.) We have the following:

Corollary 1.3. *U, V be bounded planar domains, and let $\Phi : \Sigma^n U \rightarrow \Sigma^n V$ be a proper holomorphic map. Then there is a proper holomorphic $\varphi : U \rightarrow V$ such that $\Phi = \Sigma^n \varphi$.*

Therefore, the study of boundary behavior of proper maps between equidimensional symmetric products is reduced to the study of boundary behavior of maps of the form $\Sigma^n \psi$ as defined in (1.1). Study of regularity of $\Sigma^n \psi$ is a major thrust of this paper. For k a non-negative integer or ∞ , and a domain $\Omega \subset \mathbb{C}^m$, denote by $\mathcal{A}^k(\Omega)$ the space $\mathcal{O}(\Omega) \cap \mathcal{C}^k(\bar{\Omega})$ of holomorphic functions on Ω which are C^k -smooth up to the boundary. A map is of class $\mathcal{A}^k(\Omega)$ if each component is. We prove the following result:

Theorem 1.4. *Let k be a non-negative integer or ∞ , let U, V be bounded domains in \mathbb{C} with rectifiable boundaries, and let $\varphi : U \rightarrow V$ be a holomorphic map such that $\varphi \in \mathcal{A}^{kn}(U)$. Then the induced map*

$\Sigma^n \varphi : \Sigma^n U \rightarrow \Sigma^n V$ is in $\mathcal{A}^k(\Sigma^n U)$. Further, if $\varphi : U \rightarrow V$ is a holomorphic map which is not of class $\mathcal{A}^{kn}(U)$, then the induced map is not of class $\mathcal{A}^k(\Sigma^n U)$.

Note the loss of smoothness from kn times differentiability at the boundary to k times differentiability at the boundary in going from φ to $\Sigma^n \varphi$. This is not unexpected given the serious branching behavior of the symmetrization map $\pi : \mathbb{C}^n \rightarrow \mathbb{C}^n$. It would be interesting to obtain precise information on the smoothness of the map $\Sigma^n \varphi$ when φ is of class $\mathcal{A}^r(U)$ where r is not an integer multiple of n , or for φ in Hölder classes. Combining Theorem 1.4 with classical results on boundary regularity of proper holomorphic maps in one variable, we obtain information on the boundary regularity of proper maps between symmetric products, e.g.,

Corollary 1.5. *Let $U, V \Subset \mathbb{C}$ be domains with C^∞ -smooth boundaries, and let $\Phi : \Sigma^n U \rightarrow \Sigma^n V$ be a proper holomorphic map. Then Φ extends as a C^∞ -smooth map from $\Sigma^n \bar{U}$ to $\Sigma^n \bar{V}$.*

1.3. General remarks. Topological symmetric products, under the name *unordered configuration spaces* have been studied extensively. One usually restricts attention to the “unbranched part” of such a space, i.e., one considers the quotient $(X^n \setminus \Delta)/S_n$, where Δ consists of those points (x_1, \dots, x_n) of X^n , where at least two of the coordinates x_i and x_j are the same. The fundamental group of such spaces are an important subject of study under the name “Braid Groups” (see [32].)

Complex symmetric products have many applications in the theory of complex spaces, their holomorphic mappings and correspondences (see, e.g., [41, 6].) For a compact Riemann surface C , the points of $\Sigma^n C$ are in bijection with the effective divisors on C of degree n , and the n -th Abel-Jacobi map maps $\Sigma^n C$ into the Jacobian variety of C (birationally if n is equal to the genus of C , see [5, p. 18 ff.].)

Denoting by \mathbb{D} the unit disc in the plane, the domain $\Sigma^n \mathbb{D}$ (under the name *symmetrized polydisc*, or for $n = 2$, *symmetrized bidisc*) has been studied extensively. It arises naturally in certain problems in control theory and the related spectral Nevanlinna-Pick interpolation problem (see [1], and the subsequent work deriving from it.) Subsequently, the domains $\Sigma^n \mathbb{D}$ have been studied from the point of complex geometry (see e.g. [2, 18, 19, 34, 22, 23, 3, 4] etc.) In view of Lempert’s theorem on the identity of the Carathéodory and Kobayashi metrics on convex domains, the most striking feature is the identity of these metrics on $\Sigma^2 \mathbb{D}$, though $\Sigma^2 \mathbb{D}$ cannot be exhausted by an increasing sequence of domains biholomorphic to convex domains. This however seems to be a special property of the symmetrized bidisc.

Our work here considers symmetric products of general planar domains, with particular reference to boundary regularity, and generalizes properties of symmetrized polydiscs.

The paper is organized as follows. In Section 2 below, we consider some general properties of symmetric products as a preliminary to the proof of the results stated above. We prove a formula (see (2.6)) for the induced map $\Sigma^n \varphi$ which plays a crucial role in the sequel, as well as study the branching behavior of the symmetrization mapping $\pi : \mathbb{C}^n \rightarrow \mathbb{C}^n$. Section 3 is devoted to obtaining some estimates in spaces of holomorphic functions smooth up to the boundary on an integral operator J (see (3.1).) This operator arises in the proof of Theorem 1.4, and through Theorem 1.4 in the proof of other results of this paper. The estimates are then used in the following section to prove Theorem 1.4. In Section 5, Theorem 1.1 is proved first for a model domain using a construction of a Stein neighborhood basis. Conformal mapping and a pullback argument gives the general result. Similarly, a pullback method is used to show that the boundary of a symmetric product is not Lipschitz. The last section is devoted to the proof of Theorem 1.2 and deducing Corollaries 1.3 and 1.5.

Starting with Section 3, we use the “Variable Constant” convention, so that C or C_j may denote different constants at different appearances. The convention is standard with inequalities and estimates, but here we use it in the context of equalities as well, for constants which depend only on the dimension and domain.

1.4. Acknowledgements. We would like to thank our colleagues at Bangalore for many interesting discussions. We would also like to thank Włodzimierz Zwonek and Armen Edigarian for helpful information and Łukasz Kosiński for telling us about his result on the boundary geometry of the bidisc (Lemma 5.4 below.) Debraj Chakrabarti would like to thank Diganta Borah for his invitation to visit IISER Pune, and colleagues there for useful comments and discussions on the subject of this paper. Sushil Gorai would like to

thank Stat-Math Unit, Indian Statistical Institute, Bangalore Centre for providing a congenial atmosphere for research.

We would also like to express our special gratitude to the Dean of TIFR-CAM, Prof. Mythily Ramaswamy for facilitating our collaboration by providing chauffeured cars to travel between our institutes. Without her support this paper would never have been written.

2. ELEMENTARY PROPERTIES OF SYMMETRIC PRODUCTS

2.1. The symmetrization map. Let S_n denote the symmetric group of bijections of $I_n = \{1, \dots, n\}$. Given any set X we can define an action of S_n on the n -fold cartesian product X^n by setting

$$\sigma(x_1, \dots, x_n) = (x_{\sigma(1)}, \dots, x_{\sigma(n)}). \quad (2.1)$$

Then the symmetric product $X^{(n)}$ may be identified with the quotient X^n/S_n consisting of orbits of this action. It follows that in a category where quotients by finite groups exist, we can define symmetric products. It is known that in the category of *complex spaces* (ringed spaces locally modelled on complex analytic sets, see [27, 41]) such quotients by finite groups do indeed exist (see [11, 10].) Therefore the symmetric product of a domain in \mathbb{C}^k is in general a complex space with singularities. For a profound study of symmetric products in the complex setting see [41], especially Appendix V.

If U is a domain in \mathbb{C}^k clearly the symmetric product $U^{(n)}$ sits inside the symmetric product $(\mathbb{C}^k)^{(n)}$ as an open set. Therefore, to construct $U^{(n)}$, it suffices to construct the space $(\mathbb{C}^k)^{(n)}$ as a complex space. One way to do this ([11, 10] or [41, Chaper 5, section 5]) is to consider $(\mathbb{C}^k)^{(n)}$ as an affine algebraic variety and imbed it in \mathbb{C}^N (for large N). Indeed, the coordinate ring of $(\mathbb{C}^k)^{(n)}$ is nothing but the ring of those polynomials in n variables on \mathbb{C}^k which are left invariant by the action of S_n described in the previous section. When $k = 1$, this ring is the ring of symmetric polynomials in n variables. It is well known that this ring is freely generated by the elementary symmetric polynomials. We recall that the elementary symmetric polynomials $\{\pi_k\}_{k=1}^n$ can be characterized by the fact that

$$(t - z_1)(t - z_2) \dots (t - z_n) = t^n - \pi_1(z)t^{n-1} + \dots + (-1)^n \pi_n(z), \quad (2.2)$$

and are given explicitly by

$$\pi_k(z_1, \dots, z_n) = \sum_{\substack{J \subset I_n \\ |J|=k}} \left(\prod_{j \in J} z_j \right) = \sum_{1 \leq j_1 \leq \dots \leq j_k \leq n} z_{j_1} \dots z_{j_k}.$$

In particular, the symmetrization map $\pi = (\pi_1, \dots, \pi_n) : \mathbb{C}^n \rightarrow \mathbb{C}^n$, gives a concrete realization of the quotient map $\mathbb{C}^n \rightarrow \mathbb{C}^{(n)} = \mathbb{C}^n/S_n$. We note a number of elementary properties of the map π :

(a) For $s \in \mathbb{C}^n$, a point $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ is in $\pi^{-1}(s)$, if and only if z_1, \dots, z_n are roots of the polynomial $q(s; t)$, where

$$q(s; t) = t^n - s_1 t^{n-1} + \dots + (-1)^n s_n. \quad (2.3)$$

This is an immediate consequence of (2.2).

(b) From the above, it follows that the map π is surjective, thanks to the fundamental theorem of algebra.

(c) We can define the biholomorphic map identifying \mathbb{C}^n with $\mathbb{C}^{(n)}$ in the following way. For an $s \in \mathbb{C}^n$, let (z_1, \dots, z_n) be the roots of the polynomial q of equation (2.3). Then the tuple (z_1, \dots, z_n) has no natural order, and hence may be considered to be a point in the symmetric product $\mathbb{C}^{(n)}$. This defines a map $\psi : \mathbb{C}^n \rightarrow \mathbb{C}^{(n)}$, $s \mapsto (z_1, \dots, z_n)$. This is easily seen to be a set-theoretic bijection, and using the complex structure on $\mathbb{C}^{(n)}$ one can show that this identification is biholomorphic (see [41] for details.) Similarly, for any subset U of \mathbb{C}^n , the set $\Sigma^n U$ can be identified with $U^{(n)}$.

(d) The map π is open, i.e., if $\omega \subset \mathbb{C}^n$ is open, then so is $\pi(\omega)$. This is a consequence of the classical fact that roots of a monic polynomial with complex coefficients depend continuously on the coefficients (cf. [41, Appendix V (Sec. 4)])

- (e) By the previous part, if $U \subset \mathbb{C}$ is a domain in \mathbb{C} , the set $\Sigma^n U = \pi(U^n)$ is a domain in \mathbb{C}^n . Furthermore, $\Sigma^n U$ is pseudoconvex: if ψ is a subharmonic exhaustion function of the domain U (which always exists), the function Φ given on $\Sigma^n U$ by $\Phi(s) = \sum_{j=1}^n \psi(z_j)$, where (z_1, \dots, z_n) are the roots of $q(s; t) = 0$ is well-defined and a plurisubharmonic exhaustion of $\Sigma^n U$.
- (f) The map π is proper, i.e., the inverse image of a compact set is compact under π . Indeed, if $|s| < r$, and $|t| > \max(\sqrt{nr}, 1)$, we have

$$|q(s; t)| > |t|^n - \sqrt{n}|s||t|^{n-1} > 0.$$

Consequently, the inverse image $\pi^{-1}(B(0, r))$ is contained in the ball $B(0, \max(\sqrt{nr}, 1))$, and therefore π is proper.

We note here that if $k \geq 2, n \geq 2$, the space $(\mathbb{C}^k)^{(n)}$ is *not* a smooth complex manifold. For example, the complex space $(\mathbb{C}^2)^{(2)}$ may be biholomorphically identified with the singular complex hypersurface in \mathbb{C}^5 with equation

$$4(z_3 z_5 - z_4^2) = z_2^2 z_3 - 2z_1 z_2 z_4 + z_1^2 z_5.$$

For details see [36, pp. 78-79, especially equation 4.18].

2.2. The canonical stratification. By a *partition* of a set I , we mean a collection of subsets of I which are pairwise disjoint and whose union is I . The *length* of a partition is simply the number of subsets of I in the partition. For any positive integer m , let I_m denote the m -element set $\{1, \dots, m\}$. We denote by \mathcal{P}_k^n the set of partitions of I_n of length k .

For each partition $P = \{P_1, \dots, P_k\} \in \mathcal{P}_k^n$ of I_n of length k , we define

$$S(P) = \left\{ z \in \mathbb{C}^n \mid \begin{array}{l} \text{there are } \textit{distinct} \text{ points } w_1, \dots, w_k \in \mathbb{C} \text{ such that} \\ \text{for each } \ell \in I_k, \text{ and for each } i \in P_\ell, \text{ we have } z_i = w_\ell \end{array} \right\}. \quad (2.4)$$

If we also let

$$\bar{S}(P) = \left\{ z \in \mathbb{C}^n \mid \begin{array}{l} \text{there are points } w_1, \dots, w_k \in \mathbb{C} \text{ such that for} \\ \text{each } \ell \in I_k, \text{ and for each } i \in P_\ell, \text{ we have } z_i = w_\ell \end{array} \right\},$$

(so that w_1, \dots, w_k are no longer assumed distinct) then clearly $\bar{S}(P)$ is a linear subspace of \mathbb{C}^n of dimension k with linear coordinates (w_1, \dots, w_k) . Then $S(P)$ is a dense open subset of this linear space, and the complement $\bar{S}(P) \setminus S(P)$ is an analytic set defined by $\prod_{j < k} (w_j - w_k) = 0$. Consequently $S(P)$ is a connected complex manifold of dimension k (indeed a domain of dimension k .) We also define

$$\mathbf{V}_k = \bigcup_{P \in \mathcal{P}_k^n} S(P),$$

where clearly the union is disjoint. Then \mathbf{V}_k is precisely the set of points $z \in \mathbb{C}^n$, whose coordinates (z_1, \dots, z_n) take k distinct numbers as values. Note also that each $S(P)$ is open in \mathbf{V}_k , and in fact is a connected component of \mathbf{V}_k . The following fact will be used later

Lemma 2.1. *For each $k \in I_n$, the restriction $\pi|_{\mathbf{V}_k}$ is a local biholomorphism.*

Proof. Let $z^* \in \mathbf{V}_k$, and set $s^* = \pi(z^*)$. We need to find a neighborhood D of z^* in \mathbf{V}_k such that $\pi|_D$ is a biholomorphism onto $\pi(D)$. To define D , first note that there is a partition $P \in \mathcal{P}_k^n$ such that z^* is in $S(P)$. Let w_1, \dots, w_k be the natural coordinates on $S(P)$ and suppose that the natural coordinates of z^* are (w_1^*, \dots, w_k^*) . Since the points w_1^*, \dots, w_k^* are all distinct, we can find open discs D_1, \dots, D_k in \mathbb{C} , centered at w_1^*, \dots, w_k^* , such that their closures are pairwise disjoint. We define the open subset D of $S(P)$ by setting

$$D = \{z \in S(P) : \text{for } j \in I_k, \text{ we have } w_j \in D_j\}.$$

Then D is an open neighborhood of the point z^* in $S(P)$. Let $G \subset \mathbb{C}^n$ be given by

$$G = \left\{ s \in \mathbb{C}^n : \text{no root of } q(s; t) \text{ lies in } \bigcup_{j=1}^k \text{b}D_j \right\},$$

where q is as in (2.3). Since the roots of a monic polynomial depend continuously on the coefficients, it follows that the set G is open. For the partition $P = \{P_1, \dots, P_k\}$, and $j \in I_n$, there is an $\mu(j)$ such that $j \in P_{\mu(j)}$. For $s \in G$, define for each $j \in I_n$,

$$\gamma_j(s) = \frac{1}{2\pi i |P_{\mu(j)}|} \int_{\text{b}D_{\mu(j)}} t \cdot \frac{q'(s; t)}{q(s; t)} dt,$$

where q is as in (2.3) and $q'(s; t)$ is the partial derivative

$$\frac{\partial q}{\partial t}(s; t) = nt^{n-1} - (n-1)s_1 t^{n-2} + \dots + (-1)^{n-1} s_{n-1}. \quad (2.5)$$

By differentiating with respect to $(\bar{s}_1, \dots, \bar{s}_n)$ under the integral sign, and noting that the integral on the right hand side is well defined if $s \in G$, we see that the tuple $\gamma = (\gamma_1, \dots, \gamma_n)$ defines a holomorphic map from G to \mathbb{C}^n .

Note that $\pi(D) \subset G$. For $s \in \pi(D)$, let $z \in D \subset S(P)$ be such that $\pi(z) = s$. For a given $j \in I_n$, the integrand $t \cdot \frac{q'(s; t)}{q(s; t)}$ has a pole at $t = z_j$ of order $|P_{\mu(j)}|$ in the disc $D_{\mu(j)}$. Applying the residue formula we see that $\gamma_j(s) = z_j$, i.e., $\gamma : \pi(D) \rightarrow D$ is the inverse of the map $\pi : D \rightarrow \pi(D)$. The result is proved. \square

Corollary 2.2. *The mapping $\pi|_{V_1}$ from V_1 onto $\pi(V_1)$ is a biholomorphism.*

Proof. Thanks to Lemma 2.1, we only need to show that the mapping π is injective on V_1 . The set V_1 consists of points such that all the coordinates are equal. Denoting the natural coordinate on V_1 as w , we see that the image of the point $(w, \dots, w) \in V_1$ is $\left(nw, \frac{n(n-1)}{2} w^2, \dots, w^n \right)$, which immediately shows that π is injective on V_1 . \square

2.3. The induced map $\Sigma^n \varphi$. Let $\varphi : X \rightarrow Y$ be a mapping of sets. Then we can naturally define a map $\varphi^n : X^n \rightarrow Y^n$ by setting

$$\varphi^n(x_1, \dots, x_n) = (\varphi(x_1), \dots, \varphi(x_n)).$$

The map φ^n is equivariant with respect to the action (2.1) of S_n , i.e. for each $\sigma \in S_n$, thought of as acting on X^n or Y^n , we have $\sigma \circ \varphi^n = \varphi^n \circ \sigma$. Then clearly, φ^n maps the S_n -orbits of X^n to the S_n -orbits of Y^n , and consequently induces a map $\varphi^{(n)} : X^{(n)} \rightarrow Y^{(n)}$, which we will call the *n-fold symmetric power of the map φ* . It is clear that this construction is functorial, i.e., if $\varphi : X \rightarrow Y$ and $\psi : Y \rightarrow Z$ are given maps, then $(\psi \circ \varphi)^{(n)} = \psi^{(n)} \circ \varphi^{(n)}$ as maps from $X^{(n)}$ to $Z^{(n)}$. When X and Y are subsets \mathbb{C} , and the symmetric products $X^{(n)}$ and $Y^{(n)}$ are identified with the sets $\Sigma^n X = \pi(X^n) \subset \mathbb{C}^n$ and $\Sigma^n Y = \pi(Y^n) \subset \mathbb{C}^n$, we denote the induced map $\varphi^{(n)}$ (identified with a map from $\Sigma^n X$ to $\Sigma^n Y$) by $\Sigma^n \varphi$. Consequently, $\Sigma^n \varphi$ is characterized by the equation (1.1), i.e., $(\Sigma^n \varphi)(\pi(z)) = \pi(\varphi^n(z))$, or equivalently by the commutative diagram:

$$\begin{array}{ccc} X^n & \xrightarrow{\varphi^n} & Y^n \\ \pi \downarrow & & \pi \downarrow \\ \Sigma^n X & \xrightarrow{\Sigma^n \varphi} & \Sigma^n Y \end{array}$$

We first note the following:

Lemma 2.3. *In the above diagram, $\Sigma^n \varphi$ is continuous if and only if φ is continuous.*

Proof. Assume that φ is continuous, and let ω be an open subset of $\Sigma^n Y$. We need to show that $(\Sigma^n \varphi)^{-1}(\omega)$ is open in $\Sigma^n X$. Let Ω be an open set in \mathbb{C}^n such that $\omega = \Omega \cap \Sigma^n X$. Then

$$\begin{aligned} (\Sigma^n \varphi)^{-1}(\omega) &= (\Sigma^n \varphi)^{-1}(\Omega) \cap (\Sigma^n \varphi)^{-1}(Y) \\ &= \pi((\varphi^n \circ \pi)^{-1}(\Omega)) \cap \Sigma^n X. \end{aligned}$$

Since $\varphi^n \circ \pi$ is continuous and π is open by (d) of Section 2.1, it follows that $(\Sigma^n \varphi)^{-1}(\omega)$ is open and therefore $\Sigma^n \varphi$ is continuous.

Now suppose $\Sigma^n \varphi$ is continuous. Let ψ be its restriction to $\pi(\mathbf{V}_1 \cap X^n)$. The image of ψ is contained in $\pi(\mathbf{V}_1 \cap Y^n)$. By Corollary 2.2 the map π induces a biholomorphism from \mathbf{V}_1 to $\pi(\mathbf{V}_1)$, there we have

$$\varphi^n|_{\mathbf{V}_1 \cap X^n} = \pi^{-1} \circ \psi \circ (\pi|_{\mathbf{V}_1 \cap X^n}),$$

where π^{-1} is the biholomolomorphic inverse of π from $\pi(\mathbf{V}_1)$ to \mathbf{V}_1 . Since ψ is continuous, so is $\varphi^n|_{\mathbf{V}_1 \cap X^n}$. But this last map is given as $(z, \dots, z) \mapsto (\varphi(z), \dots, \varphi(z))$ which is continuous if and only if φ is continuous. \square

Proposition 2.4. *There is a polynomial automorphism \mathfrak{P} of \mathbb{C}^n with the following property. If $U \subset \mathbb{C}$ is a domain with smooth boundary, and $\varphi : U \rightarrow V$ is a holomorphic map into a bounded domain $V \subset \mathbb{C}$, then*

$$\Sigma^n \varphi = \mathfrak{P} \circ \Psi, \quad (2.6)$$

where $\Psi : \Sigma^n U \rightarrow \mathbb{C}^n$ is the map whose m -th component function is given, for $m \in I_n$, by

$$\Psi_m(s) = \frac{1}{2\pi i} \int_{\partial U} (\varphi(t))^m \frac{q'(s; t)}{q(s; t)} dt, \quad (2.7)$$

where $s \in \Sigma^n U$, $q(s; t)$ is as in (2.3) and $q'(s; t)$ denotes the partial derivative $\frac{\partial q}{\partial t}(s; t)$ given by (2.5).

Proof. Define the power-sum map $\tau : \mathbb{C}^n \rightarrow \mathbb{C}^n$, where for $j \in I_n$, the component τ_j is the j -th power-sum polynomial:

$$\tau_j(z) = \sum_{\ell=1}^n (z_\ell)^j.$$

We claim that there exists a polynomial automorphism $\mathfrak{P} : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that

$$\pi = \mathfrak{P} \circ \tau. \quad (2.8)$$

It is known from classical algebra (cf. [31, pp. 312 ff.]) that the either of the collections $\{\tau_j\}_{j=1}^n$ or $\{\pi_j\}_{j=1}^n$ freely generates the ring of symmetric polynomials in n variables. It follows that for $j \in I_n$, there are polynomials $\mathfrak{P}_j, \mathfrak{Q}_j$ in n variables such that $\pi_j = \mathfrak{P}_j(\tau_1, \dots, \tau_n)$ and $\tau_j = \mathfrak{Q}_j(\pi_1, \dots, \pi_n)$. It is possible to write down the polynomials $\mathfrak{P}_j, \mathfrak{Q}_j$ explicitly using the Newton identities for symmetric polynomials (cf. [31, p. 323].) Define polynomial self-maps \mathfrak{P} and \mathfrak{Q} of \mathbb{C}^n by setting $\mathfrak{P} = (\mathfrak{P}_1, \dots, \mathfrak{P}_n)$ and $\mathfrak{Q} = (\mathfrak{Q}_1, \dots, \mathfrak{Q}_n)$. Then (2.8) holds and $\mathfrak{P} \circ \mathfrak{Q}$ is the identity on \mathbb{C}^n , and therefore \mathfrak{P} is a polynomial automorphism of \mathbb{C}^n .

Recall that $\mathbf{V}_n \subset \mathbb{C}^n$ is the set of points whose coordinates are n distinct complex numbers, and assume that s is a point in $\pi(\mathbf{V}_n) \cap \Sigma^n U$, i.e., the roots of $q(s; t) = 0$ are n distinct points z_1, \dots, z_n in U . Then the integrand in (2.7) is a meromorphic function on U with simple poles at the points $t = z_1, \dots, t = z_n$. We can evaluate the integral using the residue theorem, which leads to the representation

$$\begin{aligned} \Psi_m(s) &= \sum_{j=1}^n (\varphi(z_j))^m \\ &= \tau_m(\varphi(z_1), \dots, \varphi(z_n)) \\ &= \tau_m(\varphi^n(z)), \end{aligned}$$

so that we have

$$\Psi(s) = \tau(\varphi(z_1), \dots, \varphi(z_n)) = \tau(\varphi^n(z)).$$

Therefore, with \mathfrak{P} as in (2.8) we have

$$\begin{aligned}\mathfrak{P}(\Psi(s)) &= \mathfrak{P}(\tau(\varphi(z_1), \dots, \varphi(z_n))) \\ &= \pi(\varphi(z_1), \dots, \varphi(z_n)) \\ &= (\Sigma^n \varphi)(s),\end{aligned}$$

where we have used the relation (1.1) defining $\Sigma^n \varphi$. Therefore (2.6) is established if $s \in \pi(\mathbf{V}_n) \cap \Sigma^n U$. To complete the proof we note that by Lemma 2.3, $\Sigma^n \varphi$ is a continuous mapping from $\Sigma^n U$ to $\Sigma^n V$. We also claim that the map $\Psi : \Sigma^n U \rightarrow \mathbb{C}^n$ is holomorphic. Indeed, for any $j \in I_n$, by differentiation under the integral sign we have

$$\frac{\partial}{\partial \bar{s}_j} \Psi_m(s) = \frac{1}{2\pi i} \int_{\mathbf{b}U} (\varphi(t))^m \frac{\partial}{\partial \bar{s}_j} \left(\frac{q'(s; t)}{q(s; t)} \right) dt = 0. \quad (2.9)$$

Therefore, in the equation (2.6) we see that both sides extend continuously to the complement of \mathbf{V}_n , and the result follows. \square

Corollary 2.5. *Let $\varphi : U \rightarrow V$ be a mapping of planar domains. Then the induced map $\Sigma^n \varphi : \Sigma^n U \rightarrow \Sigma^k V$ is holomorphic if and only if φ is holomorphic.*

Proof. First assume that φ is holomorphic. For smoothly bounded U it was proved in course of the proof of Proposition 2.4 that $\Sigma^n \varphi$ is holomorphic. Otherwise, let $\{U_\nu\}_\nu$ be a family of bounded domains with smooth boundary such that $U_\nu \subset U_{\nu+1}$ and $\bigcup U_\nu = U$. Clearly $\Sigma^n(\varphi|_{U_\nu}) = (\Sigma^n \varphi)|_{\Sigma^n U_\nu}$, and by the previous proposition, we can write $\Sigma^n(\varphi|_{U_\nu}) = \mathfrak{P} \circ \Psi$, with Ψ as defined by (2.7) with U replaced by U_ν . Further, by differentiation under the integral sign as in (2.9) we see that Ψ and therefore $\Sigma^n(\varphi|_{U_\nu}) = (\Sigma^n \varphi)|_{\Sigma^n U_\nu}$ is holomorphic. Since this is true for each ν the result follows.

If $\Sigma^n \varphi$ is holomorphic, so is its restriction to the one-dimensional complex submanifold $\pi(\mathbf{V}_1) \cap \Sigma^n U$. But using the biholomorphism $\pi|_{\mathbf{V}_1}$, we see that this induces a holomorphic map from $\mathbf{V}_1 \cap U^n$ to $\mathbf{V}_1 \cap V^n$. Using the natural coordinate w on \mathbf{V}_1 we see that φ is itself holomorphic. \square

3. A CAUCHY-TYPE INTEGRAL

Let U be a bounded domain in the complex plane with rectifiable boundary, so that we have the Cauchy theory at our disposal. For a positive integer m we consider the function Ju on $\Sigma^n U$ defined by

$$Ju(s) = \int_{\mathbf{b}U} \frac{u(t)}{q(s; t)^m} dt \quad (3.1)$$

where q is as in (2.3). (Of course J depends on m, n but for simplicity we suppress this from the notation.) Clearly J is a linear operator from $\mathcal{C}(\mathbf{b}U)$ to $\mathcal{O}(\Sigma^n U)$, since the kernel $q(s; t)$ is holomorphic in the parameter $s \in U^n$.

For a domain $U \subset \mathbb{C}$ with smooth boundary, and an integer $k \geq 0$, let $\mathcal{A}^k(U)$ denote the space of functions $\mathcal{C}^k(\bar{U}) \cap \mathcal{O}(U)$, i.e. functions which are k times continuously differentiable on \bar{U} which are holomorphic in U . We note that $\mathcal{A}^k(U)$ is a closed subspace of the Banach space $\mathcal{C}^k(\bar{U})$, and therefore a Banach space with the norm of $\mathcal{C}^k(\bar{U})$. Let $\mathcal{A}(\Sigma^n U)$ denote the subspace of the space $\mathcal{C}(\overline{\Sigma^n U})$ of continuous functions on $\overline{\Sigma^n U}$ which are holomorphic on $\Sigma^n U$.

The following result will be used in the proof of Theorems 1.4 and 1.1.

Proposition 3.1. *For $m \geq 1$, the operator J is bounded from $\mathcal{A}^{mn-1}(U)$ to $\mathcal{A}(\Sigma^n U)$. If u is a holomorphic function on U which is not in $\mathcal{A}^{mn-1}(U)$, then Ju is not in $\mathcal{A}(\Sigma^n U)$.*

The proof will be in several steps. First, let $\mathcal{O}_{\text{sym}}(U^n)$ be the space of holomorphic functions on U^n which are symmetric in the variables (z_1, \dots, z_n) . Define a linear operator from $\mathcal{C}(\mathbf{b}U)$ to $\mathcal{O}_{\text{sym}}(U^n)$ by setting

$$(T_n u)(z) = \int_{\mathbf{b}U} \frac{u(t)}{\prod_{j=1}^n (t - z_j)^m} dt. \quad (3.2)$$

(Note that the subscript in T_n refers to the dimension of the product U^n .) The kernel function $\prod_{j=1}^n (t-z_j)^{-m}$ is holomorphic in the variable $z \in U^n$ and symmetric in the variables (z_1, \dots, z_n) , so $T_n u \in \mathcal{O}_{\text{sym}}(U^n)$. Let $H_{\text{sym}}^\infty(U^n) \subset \mathcal{O}_{\text{sym}}(U^n)$ be the Banach space of bounded holomorphic functions on U^n symmetric with respect to the variables (z_1, \dots, z_n) with the supremum norm. We first prove the following:

Lemma 3.2. *The operator T_n is bounded from $\mathcal{A}^{mn-1}(U)$ to $H_{\text{sym}}^\infty(U^n)$.*

Proof. Since $T_n u \in \mathcal{O}_{\text{sym}}(U^n)$, we only need to show that $T_n u$ is a bounded function. To do this, we use the stratification described in Section 2.2. We first show that $T_n u$ is bounded on $\mathbb{V}_1 \cap U^n$, and then show inductively that if $T_n u$ is bounded on $(\cup_{j=1}^k \mathbb{V}_j) \cap U^n$, then it is also bounded on $(\cup_{j=1}^{k+1} \mathbb{V}_j) \cap U^n$. Applying the induction step n times, we conclude that $T_n u$ is bounded on U^n .

Recall that we use C or C_j to denote a constant which depends on the integers m, n and the domain U , but not on the function u . The actual value of C or C_j at various occurrences may be different.

For the base step, let $z \in \mathbb{V}_1 \cap U^n$. Therefore, there exists $w \in U$ such that $z_1 = \dots = z_n = w$. Hence, we get from (3.2) that

$$T_n u(z) = \int_{\mathbf{b}U} \frac{u(t)}{(t-w)^{nm}} dt.$$

By the Cauchy integral formula, we obtain that

$$T_n u(z) = C \left(\frac{d}{dt} \right)^{mn-1} (u(t)) \Big|_{t=w} = C u^{(mn-1)}(w).$$

Since $u \in \mathcal{C}^{mn-1}(\overline{U})$, the function $u^{(mn-1)}$ is bounded on U . Hence, $T_n u$ is bounded on $\mathbb{V}_1 \cap U^n$.

We now use induction for the proof. Assume that $T_n u$ is bounded on $(\cup_{j=1}^k \mathbb{V}_j) \cap U^n$, and we want to show that it is bounded on $(\cup_{j=1}^{k+1} \mathbb{V}_j) \cap U^n$. Let M be the supremum of $|T_n u|$ on $(\cup_{j=1}^k \mathbb{V}_j) \cap U^n$. Since $T_n u$ is continuous, it follows that there is a neighborhood W_k of $(\cup_{j=1}^k \mathbb{V}_j) \cap U^n$ such that

$$|T_n u(z)| \leq 2M \text{ for all } z \in W_k.$$

Therefore, to complete the induction step, it is enough to show that $T_n u$ is bounded on $\mathbb{V}_{k+1} \setminus W_k$.

Let $z \in \mathbb{V}_{k+1} \cap U^n$. Then there exists a partition $Q \in \mathcal{P}_{k+1}^n$ of I_n such that $z \in S(Q) \cap U^n$, where $Q = \{Q_1, \dots, Q_{k+1}\}$ and $S(Q)$ is as defined in (2.4). We write v_1, \dots, v_{k+1} as the distinct coordinates corresponding to Q_1, \dots, Q_{k+1} in the partition Q , respectively. Therefore, again, from (3.2), we get that

$$T_n u(z) = \int_{\mathbf{b}U} \frac{u(t)}{\prod_{l=1}^{k+1} (t-v_l)^{|Q_l|m}} dt.$$

Since the integrand is a meromorphic function on U extending smoothly to $\mathbf{b}U$, we can use the residue formula to compute its contour integral:

$$T_n u(z) = \sum_{j=1}^{k+1} C_j \operatorname{res}_{t=v_j} h(t),$$

where

$$h(t) = \frac{u(t)}{\prod_{l=1}^{k+1} (t-v_l)^{|Q_l|m}}.$$

Using a classical formula (cf. [25, Proposition 4.5.6]) to compute the residue at the pole v_j of multiplicity $|Q_j| m$, we have

$$\begin{aligned} \operatorname{res}_{t=v_j} h(t) &= (|Q_j| m - 1)! \left(\frac{d}{dt} \right)^{|Q_j| m - 1} \left((t - v_j)^{|Q_j| m} \cdot h(t) \right) \Big|_{t=v_j} \\ &= (|Q_j| m - 1)! \left(\frac{d}{dt} \right)^{|Q_j| m - 1} \left(\frac{u(t)}{\prod_{l \in I_{k+1} \setminus \{j\}} (t - v_l)^{|Q_l| m}} \right) \Big|_{t=v_j}. \end{aligned}$$

Therefore,

$$T_n u(z) = \sum_{j=1}^{k+1} C_j \left(\frac{\partial}{\partial v_j} \right)^{|Q_j| m - 1} \left(\frac{u(v_j)}{\prod_{l \in I_{k+1} \setminus \{j\}} (v_j - v_l)^{|Q_l| m}} \right),$$

where the terms on the right hand side are evaluated formally. Hence, we obtain

$$|T_n u(z)| \leq C \|u\|_{\mathcal{C}^{mn-1}(\bar{U})} \cdot \sum_{j=1}^{k+1} \prod_{l \in I_{k+1} \setminus \{j\}} |v_j - v_l|^{-(|Q_l| + |Q_j|) m + 1}. \quad (3.3)$$

If $z \in (S(Q) \cap U^n) \setminus W_k$, then $z_i = v_l$ for all $i \in Q_l$, $l = 1, \dots, k+1$. Fix an r , $1 \leq r \leq k+1$. We now claim that there exists $\epsilon > 0$ such that

$$|v_r - v_l| \geq \epsilon \text{ for all } l = 1, \dots, k+1, l \neq r.$$

Suppose no such ϵ exists. Then there exists p with $1 \leq p \leq k+1$, $p \neq r$, such that for every $\epsilon > 0$, there exist $z \in (S(Q) \cap U^n) \setminus W_k$ such that

$$|v_r - v_p| < \epsilon.$$

Using the Bolzano-Weierstrass theorem, we obtain a $z^* \in (S(Q) \cap U^n) \setminus W_k$ such that for the corresponding v_j^* , we have $v_p^* = v_r^*$, which means that $z^* \in S(Q^*)$, where Q^* is a partition of length k or less. But this contradicts the fact that z^* is a limit point of a set bounded away from $\bigcup_{j=1}^k V_j$. Therefore, there exists $\epsilon > 0$ such that $|v_r - v_l| \geq \epsilon$ for all $l = 1, \dots, k+1, l \neq r$. Hence, we get from (3.3) that

$$|T_n u(z)| \leq C \|u\|_{\mathcal{C}^{mn-1}(\bar{U})} \epsilon^{-N}, \text{ for all } z \in (V_{k+1} \cap U^n) \setminus W_k,$$

where $N > 0$ is an integer.

Since we already know that $T_n u$ is bounded on W_k , we conclude that $T_n u$ is bounded on V_{k+1} . Hence, the induction step is complete. Therefore, proceeding finitely many steps, we get that $T_n u$ is a bounded function on U^n . Hence, $T_n u \in H_{\text{sym}}^\infty(U)$ for all $u \in \mathcal{A}^{mn-1}(U)$.

We therefore have shown that T_n is algebraically a linear map from $\mathcal{A}^{mn-1}(U)$ to $H_{\text{sym}}^\infty(U^n)$. To show that T_n is a bounded operator from $\mathcal{A}^{mn-1}(U)$ to $H_{\text{sym}}^\infty(U^n)$, by the closed graph theorem, it suffices to show that T_n is a closed operator. Let $\{u_\nu\} \subset \mathcal{A}^{mn-1}(U)$ be a sequence such that $\{u_\nu\}$ converges to u in $\mathcal{A}^{mn-1}(U)$ and $T_n u_\nu$ converges to v in $H_{\text{sym}}^\infty(U)$ as $\nu \rightarrow \infty$. We want to show that $v = T_n u$. Suppose that the smooth boundary $\mathbf{b}U$ is the union of M smooth closed curves, and let them be parametrized as $\gamma_k : [0, 1] \rightarrow \mathbb{C}$, where $k \in I_M$, and each γ_k is smooth. From (3.2), we have for each $z \in U^n$:

$$\begin{aligned} T_n u_\nu(z) &= \int_{\mathbf{b}U} \frac{u_\nu(t)}{\prod_{j=1}^n (t - z_j)^m} dt \\ &= \sum_{k=1}^M \int_0^1 \frac{u_\nu(\gamma_k(\tau)) \gamma_k'(\tau)}{\prod_{j=1}^n (\gamma_k(\tau) - z_j)^m} d\tau. \end{aligned}$$

Since $\{u_\nu\}$ converges to u in $\mathcal{A}^{mn-1}(U)$, a fortiori, the sequence $\{u_\nu\}$ converges uniformly to u on $\mathbf{b}U$, so the integrand in the k -th integral converges uniformly to

$$\tau \mapsto \frac{u(\gamma_k(\tau))\gamma_k'(\tau)}{\prod_{j=1}^n (\gamma_k(\tau) - z_j)^m}.$$

Therefore, taking the limit under the integral signs, we conclude that for $z \in U^n$,

$$v(z) = \lim_{\nu \rightarrow \infty} T_n u_\nu(z) = \int_{\mathbf{b}U} \frac{u(t)}{\prod_{j=1}^n (t - z_j)^m} dt = T_n u(z),$$

which completes the proof. \square

Lemma 3.2 allows us to conclude a weak version of Proposition 3.1:

Corollary 3.3. *The operator J is bounded from $\mathcal{A}^{mn-1}(U)$ to $H^\infty(\Sigma^n U)$.*

Proof. Let $\pi^* : H^\infty(\Sigma^n U) \rightarrow H_{\text{sym}}^\infty(U^n)$ be defined as $\pi^*(u) = u \circ \pi$. Then it is not difficult to see that π^* is an isometric isomorphism of Banach spaces. But it is clear that $T_n = \pi^* \circ J$. The result follows. \square

Let $\text{Lip}_\mathcal{O}(\Sigma^n U)$ denote the space of Lipschitz holomorphic functions on $\Sigma^n U$. Recall that this is a Banach space with the norm

$$\|u\|_{\text{Lip}_\mathcal{O}} = \|u\|_{\text{sup}} + \sup_{s, s' \in \Sigma^n U} \frac{|u(s) - u(s')|}{|s - s'|}.$$

Clearly, $\text{Lip}_\mathcal{O}(\Sigma^n U) \subset \mathcal{A}(\Sigma^n U)$ with continuous inclusion. We can now prove the following:

Lemma 3.4. *The operator J is bounded from $\mathcal{A}^{2mn-1}(U)$ to $\text{Lip}_\mathcal{O}(\Sigma^n U)$.*

Proof. We have, for $u \in \mathcal{A}^{2mn-1}(U)$:

$$\begin{aligned} Ju(s) - Ju(s') &= \int_{\mathbf{b}U} u(t) \left(\frac{1}{q(s;t)^m} - \frac{1}{q(s';t)^m} \right) dt \\ &= \int_{\mathbf{b}U} u(t) \frac{q(s';t)^m - q(s;t)^m}{q(s;t)^m q(s';t)^m} dt. \end{aligned} \quad (3.4)$$

Factorizing the numerator of the integrand in (3.4), we have

$$\begin{aligned} q(s;t)^m - q(s';t)^m &= (q(s';t) - q(s;t)) \cdot \sum_{k=1}^{m-1} q(s;t)^k q(s';t)^{m-1-k} \\ &= \left(\sum_{j=1}^n (-1)^j (s_j - s'_j) t^{n-j} \right) \cdot \left(\sum_{\ell=0}^{n(m-1)} P_\ell(s, s') t^\ell \right) \\ &= \sum_{j=1}^n \sum_{\ell=0}^{n(m-1)} (-1)^j (s_j - s'_j) P_\ell(s, s') t^{\ell+n-j}, \end{aligned} \quad (3.5)$$

where in the second line, we have used the expression of $q(s, t)$ and $q(s'; t)$ in the first factor, and rearranged the second factor in descending powers of t , so that P_ℓ , $\ell = 0, \dots, n(m-1)$, are polynomials on $\mathbb{C}^n \times \mathbb{C}^n$.

Hence, in view of (3.5), (3.4) reduces to

$$\begin{aligned} Ju(s) - Ju(s') &= \int_{\mathbf{b}U} u(t) \frac{\sum_{j=1}^n \sum_{\ell=0}^{n(m-1)} (-1)^j (s_j - s'_j) P_\ell(s, s') t^{\ell+n-j}}{q(s;t)^m q(s';t)^m} dt \\ &= \sum_{j=1}^n (-1)^j (s_j - s'_j) \left(\sum_{\ell=0}^{n(m-1)} P_\ell(s, s') \int_{\mathbf{b}U} \frac{t^{\ell+n-j} u(t)}{q(s;t)^m q(s';t)^m} dt \right). \end{aligned} \quad (3.6)$$

We now claim that for each j with $1 \leq j \leq n$ and each ℓ with $0 \leq \ell \leq n(m-1)$, the functions on $\Sigma^n U \times \Sigma^n U$ defined by

$$H_{j,\ell}(s, s') = \int_{\mathfrak{b}U} \frac{t^{\ell+n-j}u(t)}{q(s;t)^m q(s';t)^m} dt$$

(the integrals in (3.6)) are bounded. To estimate $H_{j,\ell}$, we consider the surjective map Π from U^{2n} to $\Sigma^n U \times \Sigma^n U$ given by $(z, z') \mapsto (\pi(z), \pi(z'))$, where $z = (z_1, \dots, z_n)$ and $z' = (z_{n+1}, \dots, z_{2n})$, where $z_j \in U$ for $j = 1, \dots, 2n$. To show that $H_{j,\ell}$ is bounded on $\Sigma^n U \times \Sigma^n U$, it is sufficient to show that $H_{j,\ell} \circ \Pi$ is bounded on U^{2n} . But for $(z, z') \in U^{2n}$ we have

$$\begin{aligned} H_{j,\ell} \circ \Pi(z, z') &= \int_{\mathfrak{b}U} \frac{t^{\ell+n-j}u(t)}{(q(\pi(z);t)q(\pi(z');t))^m} dt \\ &= \int_{\mathfrak{b}U} \frac{t^{\ell+n-j}u(t)}{\prod_{i=1}^{2n} (t - z_i)^m} dt \\ &= T_{2n}(t^{\ell+n-j}u) \end{aligned}$$

By Lemma 3.2 the operator T_{2n} is bounded from $\mathcal{A}^{2mn-1}(U)$ to $H_{\text{sym}}^\infty(U^{2n})$. We therefore have

$$\|H_{j,\ell} \circ \Pi\|_{\text{sup}} \leq C \|t^{\ell+n-j}u\|_{\mathcal{C}^{2mn-1}(\bar{U})} \leq C \|u\|_{\mathcal{C}^{2mn-1}(\bar{U})}.$$

Since the polynomial P_ℓ is bounded on $\Sigma^n U \times \Sigma^n U$, from (3.6) we get the estimate

$$|Ju(s) - Ju(s')| \leq C \|u\|_{\mathcal{C}^{2mn-1}(\bar{U})} |s - s'|,$$

which shows that Ju is Lipschitz on $\Sigma^n U$. And we also have

$$\begin{aligned} \|Ju\|_{\text{Lip}_{\mathcal{O}}(\Sigma^n U)} &= \|Ju\|_{\mathcal{C}(\bar{U})} + \sup_{s, s' \in \Sigma^n U} \frac{|Ju(s) - Ju(s')|}{|s - s'|} \\ &\leq C \|u\|_{\mathcal{C}^{mn-1}(\bar{U})} + C \|u\|_{\mathcal{C}^{2mn-1}(\bar{U})} \\ &\leq C \|u\|_{\mathcal{C}^{2mn-1}(\bar{U})}, \end{aligned}$$

which shows that the map J is bounded from $\mathcal{C}^{2mn-1}(\bar{U})$ to $\text{Lip}_{\mathcal{O}}(\Sigma^n U)$. \square

We can now complete the proof of Proposition 3.1:

Proof of Proposition 3.1. Since $\mathcal{A}(\Sigma^n U)$ is a closed subspace of the Banach space $H^\infty(\Sigma^n U)$, it is sufficient to show that for each $u \in \mathcal{A}^{mn-1}(U)$, we have $Ju \in \mathcal{A}(\Sigma^n U)$. Let $\mathcal{O}(\bar{U})$ denote the space of those functions $f \in \mathcal{O}(U)$ such that there is an open neighborhood $U_f \supset \bar{U}$ to which f extends holomorphically. It is a classical fact that for each $k \geq 0$, the space $\mathcal{O}(\bar{U})$ is dense in $\mathcal{A}^k(U)$. For $k = 0$, this is a famous theorem of Mergelyan, and the general case may be easily reduced to the case $k = 0$. Therefore we can find a sequence $\{u_\nu\}$ in $\mathcal{O}(\bar{U})$ such that $u_\nu \rightarrow u$ in the topology of $\mathcal{A}^{mn-1}(U)$ as $\nu \rightarrow \infty$. But for each u_ν , we have $Ju_\nu \in \text{Lip}_{\mathcal{O}}(\Sigma^n U)$ by Lemma 3.4. In particular, $Ju_\nu \in \mathcal{A}(\Sigma^n U)$. By continuity of J , we have $Ju_\nu \rightarrow Ju$ in the supremum norm in $H^\infty(\Sigma^n U)$. Since $\mathcal{A}(\Sigma^n U)$ is closed in $H^\infty(\Sigma^n U)$, it follows that $Ju \in \mathcal{A}(\Sigma^n U)$.

Now let $u \in \mathcal{O}(U)$, be such that $u \notin \mathcal{A}^{mn-1}(U)$. By Corollary 2.2, the map π is a biholomorphism when restricted to the complex line $\mathbb{V}_1 = \{wv | w \in \mathbb{C}\}$, where $v = (1, \dots, 1)$. As in the proof of Lemma 3.2, for $w \in U$, we can compute the value of the function $Ju(s)$ at the point $s = \pi(wv)$ on $\pi(\mathbb{V}_1)$ using the Cauchy integral formula:

$$\begin{aligned} Ju(s) &= \int_{\mathfrak{b}U} \frac{u(t)}{(t-w)^{mn}} dt \\ &= Cu^{(mn-1)}(w). \end{aligned}$$

Since $u^{(mn-1)}$ does not extend continuously to \bar{U} , it follows that $J(u)$ does not extend continuously to $\pi(\mathbb{V}_1) \cap \Sigma^n U$. It follows that $Ju \notin \mathcal{A}(\Sigma^n U)$. \square

4. PROOF OF THEOREM 1.4

When $k = 0$, applying Lemma 2.3 to the continuous map $\varphi : \bar{U} \rightarrow \bar{V}$, we see that $\Sigma^n \varphi$ is continuous from $\Sigma^n \bar{U}$ to $\Sigma^n \bar{V}$. Further by Corollary 2.5, the map $\Sigma^n \varphi$ is holomorphic from $\Sigma^n U$ to $\Sigma^n V$, which proves Theorem 1.4 when $k = 0$.

Therefore, we may assume $k \geq 1$. In view of (2.6), the regularity of the map $\Sigma^n \varphi$ is same as that of the mapping Ψ . We therefore want to compute the first k derivatives of the function Ψ_m and show that they extend continuously to the boundary of $\Sigma^n U$. In view of the representation (2.7) of the components Ψ_m of Ψ , the conclusion of Theorem 1.4 follows from the following lemma:

Lemma 4.1. *For $k \geq 1$, and a function $g \in \mathcal{A}^1(U)$, let the function G be defined by*

$$G(s) = \int_{\mathbf{b}U} g(t) \frac{q'(s;t)}{q(s;t)} dt, \quad (4.1)$$

then $G \in \mathcal{A}^k(\Sigma^n U)$ if and only if $g \in \mathcal{A}^{kn}(U)$.

Proof. By the Whitney extension theorem it is sufficient to show that for each multi-index α , with $|\alpha| \leq k$, the derivative $\partial^\alpha G(s)$ is in $\mathcal{A}(\Sigma^n U)$. Again, in this proof we let C denote a constant which only depends on the multi-index α . The value of C in different occurrences may be different. If $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multi-index, denote by ∂^α the complex partial derivative

$$\left(\frac{\partial}{\partial s_1} \right)^{\alpha_1} \left(\frac{\partial}{\partial s_2} \right)^{\alpha_2} \cdots \left(\frac{\partial}{\partial s_n} \right)^{\alpha_n},$$

acting on holomorphic functions of n complex variables (s_1, \dots, s_n) . We now compute $\partial^\alpha G$. If $t \in \mathbf{b}U$, the function $\frac{q'(s;t)}{q(s;t)}$ is holomorphic in $s \in \Sigma^n U$. Hence, for $1 \leq j \leq n$, we can differentiate (4.1) with respect to s_j under the integral sign to get:

$$\frac{\partial}{\partial s_j} G(s_1, \dots, s_n) = \int_{\mathbf{b}U} g(t) \frac{\partial}{\partial s_j} \left(\frac{q'(s;t)}{q(s;t)} \right) dt.$$

After choosing a locally defined branch of the logarithm of q , we have

$$\frac{\partial}{\partial s_j} \left(\frac{q'(s;t)}{q(s;t)} \right) = \frac{\partial}{\partial s_j} \left(\frac{\partial}{\partial t} \log q(s;t) \right) = \frac{\partial}{\partial t} \left(\frac{\partial}{\partial s_j} \log q(s;t) \right) = \frac{\partial}{\partial t} \left(\frac{q_j(s;t)}{q(s;t)} \right), \quad (4.2)$$

where

$$q_j(s;t) = \frac{\partial q(s;t)}{\partial s_j} = (-1)^j t^{n-j}, \quad (4.3)$$

since $q(s;t) = t^n - s_1 t^{n-1} + \cdots + (-1)^n s_n$. Since the first and last expressions in (4.2) are globally defined rational functions, they are equal for all $s \in \Sigma^n U$ and all $t \in U$. Therefore,

$$\begin{aligned} \frac{\partial}{\partial s_j} G(s_1, \dots, s_n) &= \int_{\mathbf{b}U} g(t) \frac{\partial}{\partial t} \left(\frac{q_j(s;t)}{q(s;t)} \right) dt \\ &= \int_{\mathbf{b}U} \left\{ \frac{\partial}{\partial t} \left(g(t) \frac{q_j(s;t)}{q(s;t)} \right) - g'(t) \frac{q_j(s;t)}{q(s;t)} \right\} dt \\ &= - \int_{\mathbf{b}U} g'(t) \frac{q_j(s;t)}{q(s;t)} dt \\ &= (-1)^{n-j+1} \int_{\mathbf{b}U} \frac{t^{n-j} g'(t)}{q(s;t)} dt \end{aligned} \quad (4.4)$$

since the integral of the first term in the second line vanishes. By repeated differentiation, and using (4.3) we have for each non-zero multi-index $\beta = (\beta_1, \dots, \beta_n)$ that

$$\partial^\beta \left(\frac{1}{q(s;t)} \right) = C_\beta \cdot \frac{t^{\sum_{k=1}^n (n-k)\beta_k}}{q(s;t)^{|\beta|+1}},$$

where C_β is a constant which depends only on the multi-index β .

For a multi-index $\alpha \neq 0$, there exists j with $1 \leq j \leq n$, such that $\alpha_j > 0$. Let $\beta = \alpha - e^j$, where e^j is a unit multi-index with 1 in the j -th position and 0 everywhere else. Again, differentiating under the integral sign in the equation (4.4), we have:

$$\begin{aligned} \partial^\alpha G(s) &= (-1)^{n-j+1} \int_{\mathfrak{b}U} t^{n-j} g'(t) \cdot \partial^\beta \left(\frac{1}{q(s;t)} \right) dt \\ &= (-1)^{n-j+1} \cdot C_\beta \int_{\mathfrak{b}U} t^{n-j+\sum_{k=1}^n (n-k)\beta_k} g'(t) \cdot \frac{1}{q(s;t)^{|\beta|+1}} dt \\ &= C \int_{\mathfrak{b}U} g'(t) \frac{t^\gamma}{q(s;t)^{|\alpha|}} dt, \end{aligned} \quad (4.5)$$

where C is a constant depending on the multi-index α alone and γ is given by

$$\gamma = \sum_{\ell=1}^n (n-\ell)\alpha_\ell. \quad (4.6)$$

Then, in the notation of Proposition 3.1, we can write, with $m = |\alpha|$, that:

$$\partial^\alpha G(s) = J(t^\gamma g').$$

It now follows from Proposition 3.1 that $\partial^\alpha G \in \mathcal{A}(\Sigma^n U)$ for $|\alpha| \leq k$ if and only if $t^\gamma g'$ is of class $\mathcal{A}^{kn-1}(U)$. Applying the Whitney Extension theorem, this is equivalent to G being of class $\mathcal{A}^k(U)$, and note that this also holds if $k = \infty$. \square

5. PROOF OF THEOREM 1.1

In the proof of Theorem 1.1 we use the following result [20, 15], which can be proved either by L^2 methods, or by integral representations.

Result 5.1 (Dufresnoy [20]; also see [15]). *Let Γ be a compact set in \mathbb{C}^n , and suppose that there is a family of open pseudoconvex neighborhoods $\{\Omega_\nu\}_{\nu=1}^\infty$ of Γ such that*

(1) *There is a $\theta > 0$ such that for $0 < \epsilon < \frac{1}{2}$, there is a ν such that*

$$\Gamma(\epsilon^\theta) \subset \Omega_\nu \subset \Gamma(\epsilon)$$

where $\Gamma(\delta)$ denotes the δ -neighborhood

$$\Gamma(\delta) = \{z \in \mathbb{C}^n \mid \text{dist}(z, \Gamma) < \delta\}.$$

(2) *For each μ there is a ν_0 such that if $\nu > \nu_0$, the set $\overline{\Omega_\nu}$ is holomorphically convex in Ω_μ .*

Then given a form $g \in \mathcal{C}_{p,q}^\infty(\Gamma)$ with $\bar{\partial}g = 0$, there is a $u \in \mathcal{C}_{p,q-1}^\infty(\Gamma)$ such that $\bar{\partial}u = g$.

Note that by condition (2), it follows that we have

$$\bigcap_{\nu=1}^\infty \Omega_\nu = \Gamma.$$

By definition, a \mathcal{C}^∞ -function on the set Γ is a function admitting a \mathcal{C}^∞ extension to a neighborhood.

5.1. Model case. Let $D(z, r)$ denote the open disc in the plane with centre at $z \in \mathbb{C}$ and radius $r > 0$, and let $\mathbb{D} = D(0, 1)$ be the open unit disc. A domain V in the plane is said to be a *domain with circular boundary* if there is a nonnegative integer M such that V can be represented as

$$V = \mathbb{D} \setminus \left(\bigcup_{j=1}^M \overline{D(z_j, r_j)} \right), \quad (5.1)$$

where the M closed discs $\{\overline{D}(z_j, r_j)\}_{j=1}^M$ are pairwise disjoint and are all contained in the open unit disc \mathbb{D} . The domain V is a *semi-algebraic* subset of $\mathbb{C} = \mathbb{R}^2$ (see [9, Chapter 2] for more information on semi-algebraic subsets of \mathbb{R}^n , which formally correspond to affine algebraic varieties in \mathbb{C}^n .) “Semi-algebraic” means that the set V is defined in \mathbb{R}^2 by a finite system of polynomial equations and inequalities. Setting $z_j = x_j + iy_j$, it is easy to write down the $M + 1$ polynomial inequalities that define V : the inequality $x^2 + y^2 < 1$ along with the M inequalities $(x - x_j)^2 + (y - y_j)^2 > r_j^2$, where $j \in I_M$. A mapping between semi-algebraic sets is said to be semi-algebraic if its graph is semi-algebraic. We will need the following facts (see [9]):

- (a) The class of semialgebraic subsets is closed under various natural operations. Interiors, closures, projections of semi-algebraic sets are semi-algebraic. The image of a semi-algebraic subset under a semi-algebraic mapping is semi-algebraic. In particular, if V is any semi-algebraic subset of $\mathbb{C} = \mathbb{R}^2$, we see that the symmetric product $\Sigma^n V \subset \mathbb{C}^n = \mathbb{R}^{2n}$ is semi-algebraic for each n . This follows since V^n is clearly semi-algebraic, and the mapping $\pi : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ is polynomial, and in particular semi-algebraic, so $\Sigma^n V = \pi(V^n)$ is semi-algebraic. Similarly, $\overline{\Sigma^n V} = \Sigma^n(\overline{V}) = \pi(\overline{V}^n)$ is semi-algebraic.
- (b) A *semi-algebraic function* is defined to be a real valued semi-algebraic mapping. The semi-algebraic functions on a semi-algebraic set are closed under the standard operations of taking sums, products, negatives etc, as well as under taking maxima or minima of finitely many such functions.
- (c) If A is a non-empty semi-algebraic subset of \mathbb{R}^N then $x \mapsto \text{dist}(x, A)$ is a continuous semi-algebraic function on \mathbb{R}^n , which vanishes precisely on the closure of A .
- (d) (Łojasiewicz inequality.) Let B be a compact semi-algebraic subset of \mathbb{R}^N and let $f, g : B \rightarrow \mathbb{R}$ be two continuous semi-algebraic functions such that $f^{-1}(0) \subset g^{-1}(0)$. Then there is an integer $P > 0$ and a constant $C \in \mathbb{R}$ such that $|g|^P \leq C|f|$ on B .

We can now prove a special case of Theorem 1.1:

Proposition 5.2. *If V is a domain with circular boundary, then on $\Sigma^n V$ the $\bar{\partial}$ -problem is globally regular.*

Proof. We will verify that $\Gamma = \overline{\Sigma^n V}$ satisfies the hypotheses of Result 5.1 quoted at the beginning of this section.

Representing V as in (5.1), consider the function ψ defined in a neighborhood of \overline{V} by setting

$$\psi(z) = \max \left(|z|, \max_{j=1, \dots, M} \frac{r_j}{|z - z_j|} \right) - 1 \quad (5.2)$$

Observe that $\psi < 0$ is precisely the set V , and in a neighborhood of \overline{V} the function ψ is continuous, subharmonic and semi-algebraic. Define $V_\nu = \{z \in \mathbb{C} : \psi(z) < \frac{1}{\nu}\}$, and set $\Omega_\nu = \Sigma^n(V_\nu) = \pi((V_\nu)^n)$, so that $\{\Omega_\nu\}$ is a system of pseudoconvex neighborhoods of Γ . We need to verify the two conditions of Result 5.1.

Define real valued functions f, g as follows. Let s be in a neighborhood of Γ , and let z_1, \dots, z_n be the roots of the equation $q(s; t) = 0$, with q as in (2.3), so that $\pi(z_1, \dots, z_n) = s$. Define $f(s) = \max_{j \in I_n} \psi(z_j, V)$. It is straightforward to verify that this is a well-defined continuous semi-algebraic plurisubharmonic function on a neighborhood B of $\overline{\Sigma^n V}$. After shrinking, the neighborhood B is compact and semi-algebraic. We let g be the function on B given by $g(s) = \text{dist}(s, \Sigma^n V)$, where on the right hand side the distance in \mathbb{C}^n is meant. Observe that $f^{-1}(0) = g^{-1}(0) = \overline{\Sigma^n V}$, the level set $\{s \in B : f(s) < \frac{1}{\nu}\}$ is precisely the set Ω_ν and the level set $\{s \in B : g(s) < \epsilon\}$ is the ϵ -neighborhood $\Gamma(\epsilon)$ of $\Gamma = \overline{\Sigma^n V}$.

Therefore, applying the Łojasiewicz inequality twice we see that for $s \in B$, we have

$$C_1 f(s)^{N_1} \leq g(s) \leq C_2 f(s)^{\frac{1}{N_2}}, \quad (5.3)$$

where N_1, N_2 are positive integers and C_1, C_2 are positive constants. Given small $\epsilon > 0$, choose the integer ν so that

$$\frac{1}{2} \left(\frac{\epsilon}{C_2} \right)^{N_2} < \frac{1}{\nu} < \left(\frac{\epsilon}{C_2} \right)^{N_2}.$$

Such ν exists for small ϵ . Also, let θ be so large that for all small enough η , we have

$$\frac{1}{C_1^{\frac{1}{N_1}}} \eta^{\frac{\theta}{N_1}} < \frac{1}{2C_2^{N_2}} \eta^{N_2}.$$

Let $s \in \Omega_\nu$ so that $f(s) < \frac{1}{\nu}$. Then we have by the right half of (5.3):

$$g(s) < C_2 f(s)^{\frac{1}{N_2}} < \epsilon.$$

Also if $s \in \Gamma(\epsilon^\theta)$, i.e., $g(s) < \epsilon^\theta$, we have $f(s) < \frac{1}{\nu}$, so that $\Gamma(\epsilon^\theta) \subset \Omega_\nu \subset \Gamma(\epsilon)$. This verifies the first condition.

Let $\nu > \mu$. Note that the domain Ω_μ is identical to $\{f < \frac{1}{\mu}\}$ and the subdomain Ω_ν is similarly given by $\{f < \frac{1}{\nu}\}$. Since f is plurisubharmonic, it follows that the plurisubharmonic hull of $\overline{\Omega}_\nu$ in Ω_μ is itself, i.e., $\overline{\Omega}_\nu$ is plurisubharmonically convex in Ω_μ . Since Ω_μ is pseudoconvex, the notions of plurisubharmonic convexity and holomorphic convexity coincide on Ω_μ (see [28, Theorem 4.3.4].) This establishes the second condition of Result 5.1, and therefore the the proposition. \square

Remarks: (1) In fact the closed symmetrized polydisc $\Sigma^n \overline{\mathbb{D}}$ is polynomially convex in \mathbb{C}^n (see [39, Theorem 1.6.24].)

(2) The method of Proposition 5.2 may also be used to show that the $\bar{\partial}$ -problem is globally regular on $\Sigma^n V$, when V is a *subanalytic* domain in \mathbb{C} (For definition and basic properties of subanalytic sets, see [8].) We need to replace the subharmonic function ψ of (5.2) by the function $\tilde{\psi}(z) = \text{dist}(z, V)$, where the usual distance in \mathbb{C} is meant. The function $\tilde{\psi}$ is not subharmonic in general, but if we define f as in the proof above setting $f(s) = \max_{j \in I_n} \tilde{\psi}(z_j, V)$, it is not difficult to see that the sublevel sets are still pseudoconvex, and an argument parallel to the one given above, using a subanalytic version of the Łojasiewicz inequality completes the proof.

5.2. Proof of Theorem 1.1. For $q \geq 1$, let $f \in \mathcal{C}_{p,q}^\infty(\overline{\Sigma^n U})$ be a (p, q) -form with $\bar{\partial}f = 0$. We need to show that the $\bar{\partial}$ -problem

$$\bar{\partial}u = f \quad \text{such that} \quad u \in \mathcal{C}_{p,q-1}^\infty(\overline{\Sigma^n U}) \quad (5.4)$$

has a solution u smooth up to the boundary. By well-known classical results in complex analysis of one variable (cf. [17, Theorem 7.9] and [40, Theorem IX.35, p. 424]), since $U \Subset \mathbb{C}$ with \mathcal{C}^∞ -smooth boundary, there exists a biholomorphic map φ from a domain V with circular boundary to U which extends to a \mathcal{C}^∞ diffeomorphism from \overline{V} to \overline{U} . From Theorem 1.4, it follows that $\Sigma^n \varphi : \Sigma^n V \rightarrow \Sigma^n U$ extends as a \mathcal{C}^∞ map up to $\overline{\Sigma^n V}$. Since φ is a biholomorphism, the same argument applied to φ^{-1} shows that $\Sigma^n \varphi^{-1} : \Sigma^n U \rightarrow \Sigma^n V$ is holomorphic map that extends smoothly from $\overline{\Sigma^n U}$ to $\overline{\Sigma^n V}$. We observe that $(\Sigma^n \varphi)^{-1} = \Sigma^n \varphi^{-1}$ on $\overline{\Sigma^n U}$, so that $\Sigma^n \varphi$ is a biholomorphism.

The pull back $(\Sigma^n \varphi)^* f$ of f by $\Sigma^n \varphi$, is therefore a form in $\mathcal{C}_{p,q}^\infty(\overline{\Sigma^n V})$. Denoting $g = (\Sigma^n \varphi)^* f$, and noting that $\bar{\partial}g = \bar{\partial}(\Sigma^n \varphi)^* f = (\Sigma^n \varphi)^*(\bar{\partial}f) = 0$, we consider the $\bar{\partial}$ -problem $\bar{\partial}v = g$ on $\Sigma^n V$. By Proposition 5.2 we obtain that the $\bar{\partial}$ -problem is globally regular on $\Sigma^n V$, and therefore, there exists a $v \in \mathcal{C}_{p,q-1}^\infty(\overline{\Sigma^n V})$ such that $\bar{\partial}v = g$. We now let $u = (\Sigma^n \varphi^{-1})^* v$. This is a solution of (5.4) that is in $\mathcal{C}_{p,q-1}^\infty(\overline{\Sigma^n U})$. This completes the proof of Theorem 1.1.

5.3. Non-Lipschitz nature of the boundary of a symmetric product. As a consequence of Theorem 1.4, and the use of the model (5.1) we can show that the boundary of $\Sigma^n U$ is not Lipschitz:

Proposition 5.3. *Let $U \Subset \mathbb{C}$ be a domain with \mathcal{C}^∞ boundary. For $n \geq 2$, the symmetric product $\Sigma^n U$ does not have Lipschitz boundary.*

We begin with a special case of this proposition, which was shown to us by Łukasz Kosiński:

Lemma 5.4 ([30]). *Proposition 5.3 holds for $\Sigma^2 \mathbb{D}$. In fact near the point $(2, 1) \in \mathfrak{b}\Sigma^2 \mathbb{D}$ the boundary fails to be Lipschitz.*

Proof. It is well-known (see [4]) that we can represent the domain $\Sigma^2\mathbb{D}$ as

$$\left\{ (s_1, s_2) \in \mathbb{C}^2 : |s_1 - \bar{s}_1 s_2| + |s_2|^2 < 1 \right\}. \quad (5.5)$$

We will show that there is no four dimensional cone with vertex at $(2, 1) \in \mathbf{b}\Sigma^2\mathbb{D}$ inside the symmetrized bidisc. Let ℓ be a parametrized line segment contained in the domain $\Sigma^2\mathbb{D}$ passing through the point $(2, 1)$, given by $\ell(t) = (2 - bt, 1 - at)$, where $t \in (0, r)$, $r > 0$, and $a, b \in \mathbb{C}$. In view of (5.5), we get that

$$|2 - bt - (2 - \bar{b}t)(1 - at)| + |1 - at|^2 < 1.$$

Computing the above expression we have

$$|-2it \operatorname{Im} b + 2at - a\bar{b}t^2| - 2t \operatorname{Re} a + |a|^2 t^2 < 0.$$

Dividing the above expression by t and then taking limit as $t \rightarrow 0^+$ we get $|a - i \operatorname{Im} b| \leq \operatorname{Re} a$, which is possible only when $\operatorname{Im} a = \operatorname{Im} b$. This forces any cone inside $\Sigma^2\mathbb{D}$ with vertex at $(2, 1)$ to be at most of dimension three, and consequently, the domain cannot be Lipschitz near $(2, 1)$. \square

Now let V be a domain with circular boundary, represented as in (5.1). We can strengthen the above lemma in the following way:

Lemma 5.5. *For $n \geq 2$, the domain $\Sigma^n V$ is not Lipschitz.*

Proof. Consider the polynomial map Φ from $\mathbb{C}^2 \times \mathbb{C}^{n-2}$ to \mathbb{C}^n given in the following way. For $s \in \mathbb{C}^2$, let $(z_1, z_2) \in \mathbb{C}^2$ be such that $\pi^2(z_1, z_2) = s$, where $\pi^2 : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is the symmetrization map. Similarly for $\sigma \in \mathbb{C}^{n-2}$, let $(z_3, \dots, z_n) \in \mathbb{C}^{n-2}$ be such that $\pi^{n-2}(z_3, \dots, z_n) = \sigma$, where again $\pi^{n-2} : \mathbb{C}^{n-2} \rightarrow \mathbb{C}^{n-2}$ is the symmetrization map on \mathbb{C}^{n-2} . We define $\Phi(s, \sigma) = \pi^n(z_1, \dots, z_n)$, where π^n is the symmetrization map on \mathbb{C}^n . A calculation shows that if we set $\sigma_0 = 1$, and $\sigma_{-1} = \sigma_{n-1} = \sigma_n = 0$, the components of the map $\Phi = (\Phi_1, \dots, \Phi_n)$ are given by

$$\Phi_k(s, \sigma) = s_2 \sigma_{k-2} + s_1 \sigma_{k-1} + \sigma_k.$$

Taking derivatives, we see that the Jacobian matrix of Φ is given by the $n \times n$ matrix

$$\Phi'(s, \sigma) = \begin{pmatrix} \sigma_0 & \sigma_{-1} & 1 & 0 & \dots & \dots & 0 \\ \sigma_1 & \sigma_0 & s_1 & 1 & \dots & \dots & 0 \\ \sigma_2 & \sigma_1 & s_2 & s_1 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \sigma_{n-3} & \sigma_{n-2} & 0 & \dots & s_2 & s_1 & 1 \\ \sigma_{n-2} & \sigma_{n-3} & 0 & \dots & \dots & s_2 & s_1 \\ \sigma_{n-1} & \sigma_{n-2} & 0 & 0 & \dots & 0 & s_2 \end{pmatrix}$$

Without loss of generality, we can assume that $0 \in V$. If this is not already the case, we can apply a holomorphic automorphism of the unit disc mapping some point $p \in V$ to 0, and replace V by the image under this automorphism which is still a domain with circular boundary.

Note that then the origin of \mathbb{C}^{n-2} is an interior point of the domain $\Sigma^{n-2}V$. It is clear that $\Phi(\Sigma^2V \times \Sigma^{n-2}V) = \Sigma^n V$.

Let $P = (2, 1) \times 0 \in \mathbb{C}^2 \times \mathbb{C}^{n-2}$. Since $(2, 1) \in \mathbf{b}\Sigma^2V$ and $0 \in \Sigma^{n-2}V$, it is clear that $P \in \mathbf{b}(\Sigma^2V \times \Sigma^{n-2}V)$, and therefore $\Phi(P) \in \mathbf{b}\Sigma^n V$. By the previous lemma, P is a non-Lipschitz point of the boundary of $\Sigma^2V \times \Sigma^{n-2}V$. Therefore, it suffices to show that at the point P the map Φ is a local biholomorphism.

Noting that we have $\sigma_0 = 1$, and $\sigma_{-1} = \sigma_{n-1} = \sigma_n = 0$, and substituting the coordinates of P as $\sigma = (\sigma_1, \dots, \sigma_{n-2}) = 0$, and $(s_1, s_2) = (2, 1)$, we see that $\Phi'(P) = \Phi'((2, 1), 0)$ is an upper triangular matrix with main diagonal $(1, 1, 1, \dots, 1)$. Therefore we have that $\det \Phi'(P) = 1$, and therefore Φ is a local biholomorphism in a neighborhood of P . \square

Proof of Proposition 5.3. We proceed as in the proof of Theorem 1.1. Let V be a domain with circular boundary such that there is a biholomorphic map $\varphi : V \rightarrow U$ which extends as a \mathcal{C}^∞ diffeomorphism from \overline{V} to \overline{U} . Then by Theorem 1.4, we have an induced diffeomorphism $\Sigma^n \varphi : \Sigma^n \overline{V} \rightarrow \Sigma^n \overline{U}$, which is biholomorphic in the interior. Consequently we obtain that the boundary of $\Sigma^n U$ is diffeomorphic to the boundary of $\Sigma^n V$, and the result follows since the boundary of $\Sigma^n V$ is not Lipschitz by the previous lemma. \square

6. PROOF OF THEOREM 1.2

We begin with the following lemma:

Lemma 6.1. *Let $m \geq 1$ and let $\Omega \subset \mathbb{C}^m$ and $V \subset \mathbb{C}$ be domains, and let $\varphi : \Omega \rightarrow \mathbb{C}^n$ be a holomorphic mapping whose image lies in $\mathfrak{b}(\Sigma^n V)$. Then there is a point $c \in \mathfrak{b}V$ such that for all $z \in \Omega$ we have $q(\varphi(z); c) = 0$, where q is as in (2.3).*

Proof. We use the notation of Section 2.2. Since \mathbb{C}^n is the disjoint union of the sets $\pi(\mathbf{V}_k)$, where $k \in I_n$, it follows that at least one of the sets $\varphi^{-1}(\pi(\mathbf{V}_k)) \subset \Omega$ has an interior. Therefore there is a $k \in I_n$ and a p in Ω such that there is a neighborhood of p which is mapped by φ into $\pi(\mathbf{V}_k)$. Since π is a local biholomorphism from \mathbf{V}_k onto $\pi(\mathbf{V}_k)$, we see that there is a holomorphic map ψ from a neighborhood of $\varphi(p)$ in $\pi(\mathbf{V}_k)$ to \mathbf{V}_k such that $\pi \circ \psi$ is the identity in a neighborhood of $\varphi(p)$ in $\pi(\mathbf{V}_k)$. Let $\Phi = \psi \circ \varphi$. Then Φ is a map from a neighborhood ω of p in \mathbb{C}^m such that $\pi \circ \Phi = \varphi$ on ω . It follows that the image of Φ lies in $\mathfrak{b}V^n$. For $j \in I_n$, denote by ω_j the set of points $\Phi_j^{-1}(\mathfrak{b}V)$, i.e., the points in ω mapped by the j -th component of Φ to $\mathfrak{b}V$. Since the image of Φ lies in $\mathfrak{b}V^n$, it follows that $\bigcup_{j=1}^n \omega_j = \omega$. Therefore, there is at least one $j \in I_n$ such that ω_j has an interior, which by the open mapping theorem implies that Φ_j is a constant, whose value c is in $\mathfrak{b}V$. For a fixed $s \in \mathbb{C}^n$ the roots of the equation $q(s; t) = 0$ are the coordinates of the points in $\pi^{-1}(s)$. It follows therefore that $q(\varphi(z); c) = 0$ if $z \in \omega$. Since the function $z \mapsto q(\varphi(z); c)$ is holomorphic on Ω , the result follows by the identity theorem. \square

Proof of Theorem 1.2. We proceed along the lines of the proof of the Remmert-Stein theorems (cf. [35, pp. 71–77].) Consider a sequence $\{w^\nu\} \subset U_n$ such that $w^\nu \rightarrow w^0 \in \mathfrak{b}U_n$. Denote by G_{n-1} the domain $U_1 \times \cdots \times U_{n-1}$ in \mathbb{C}^{n-1} , and define the maps φ^ν on G_{n-1} by

$$\varphi^\nu('z) = f('z, w^\nu), \quad (6.1)$$

where $'z = (z_1, \dots, z_{n-1}) \in G_{n-1}$. Since f is bounded, the sequence $\{\varphi^\nu\}$ is uniformly bounded. Hence, by Montel's theorem, there exists a subsequence $\{\nu_k\}_{k=1}^\infty$ such that φ^{ν_k} converges uniformly on compact subsets of G_{n-1} . Denote the limit by $\varphi : G_{n-1} \rightarrow \mathbb{C}^n$. Since f is a proper holomorphic map, $\varphi(G_{n-1}) \subset \mathfrak{b}\Sigma^n V$. Using Lemma 6.1 we get that there is a constant $c \in \mathfrak{b}V$ such that on G_{n-1} :

$$q(\varphi; c) = c^n - c^{n-1}\varphi_1 + \cdots + (-1)^n \varphi_n \equiv 0. \quad (6.2)$$

Differentiating (6.2) with respect to z_j for each $j \in I_{n-1}$ gives rise to the following system of equations:

$$\sum_{k=1}^n (-1)^k c^{n-k} \frac{\partial \varphi_k}{\partial z_j} = 0, \quad j \in I_{n-1}. \quad (6.3)$$

Since $f : G \rightarrow \Sigma^n V$ is a proper holomorphic map, the complex Jacobian determinant $\det(f')$ of f does not vanish identically. Therefore, there is a $\mu \in I_n$ such the $(n-1) \times (n-1)$ minor of the Jacobian determinant obtained by omitting the μ -th column is not identically zero on G . We consider the following system of linear equations in the $n-1$ variables $(-1)^k c^{n-k}$, $k \in I_n \setminus \{\mu\}$, obtained by moving the μ -th column of (6.3) to the right hand side:

$$\sum_{k \in I_n \setminus \{\mu\}} (-1)^k c^{n-k} \frac{\partial \varphi_k}{\partial z_j} = (-1)^{\mu+1} c^{n-\mu} \frac{\partial \varphi_\mu}{\partial z_j}, \quad j \in I_{n-1}. \quad (6.4)$$

Let B be an $(n-1) \times n$ matrix, and let $k \in I_n \setminus \{\mu\}$. Denote by $\Delta_k(B)$ the determinant of the $(n-1) \times (n-1)$ matrix obtained by removing the μ -th column of B and replacing it by the k -th column. Denote also by $\Delta_\mu(B)$ the determinant of the $(n-1) \times (n-1)$ matrix obtained by removing the μ -th column of B . Notice that each of the functions Δ_j is a polynomial in the entries of the matrix B .

We now solve the system (6.3) by Cramer's rule to obtain:

$$(-1)^k c^{n-k} = (-1)^\mu c^{n-\mu} \frac{\Delta_k(\mathbf{D}_{n-1}\varphi)}{\Delta_\mu(\mathbf{D}_{n-1}\varphi)}, \quad (6.5)$$

where $\mathbf{D}_{n-1}\varphi$ is the $(n-1) \times n$ matrix $\left(\frac{\partial \varphi_k}{\partial z_j} \right)_{j \in I_{n-1}, k \in I_n}$. Note that by the choice of μ , the denominator $\Delta_\mu(\mathbf{D}_{n-1}\varphi)$ does not vanish identically on G_{n-1} .

We now consider two cases. If $\mu \neq 1$, we let $k = \mu - 1$ in (6.5). Then we have on G_{n-1}

$$\Delta_{\mu-1}(\mathbf{D}_{n-1}\varphi) = -c \cdot \Delta_\mu(\mathbf{D}_{n-1}\varphi). \quad (6.6)$$

Let $\Psi_k(z) = \Delta_k(\mathbf{D}_{n-1}\varphi(z))$. Then Ψ_k is a holomorphic function on G_{n-1} . Note that Ψ_k depends on the choice of the sequence w^ν as well as the subsequence ν_k (since φ depends on these choices.) We have $\Psi_{k-1}(z) = -c\Psi_k(z)$ for all $z \in G_{n-1}$. Differentiating with respect to z_j and eliminating c we obtain the relation

$$\Psi_{\mu-1}(z) \frac{\partial \Psi_\mu}{\partial z_j}(z) - \Psi_\mu(z) \frac{\partial \Psi_{\mu-1}}{\partial z_j}(z) = 0 \quad (6.7)$$

for all $z \in G_{n-1}$. Denote by $\mathbf{D}_{n-1}^2\varphi$ the collection of second derivatives of the components $(\varphi_1, \dots, \varphi_n)$ of φ with respect to the $(n-1)$ coordinate variables z_1, \dots, z_{n-1} . From the chain rule, there is clearly a polynomial E_{jk} such that

$$\begin{aligned} \frac{\partial \Psi_k}{\partial z_j}(z) &= \frac{\partial}{\partial z_j} \Delta_k(\mathbf{D}_{n-1}\varphi(z)) \\ &= E_{jk}(\mathbf{D}_{n-1}\varphi(z); \mathbf{D}_{n-1}^2\varphi(z)). \end{aligned}$$

Therefore, if H_j is the polynomial $\Psi_{\mu-1}E_{j\mu} - \Psi_\mu E_{j-1,\mu}$ then (6.7) may be rewritten as

$$H_j(\mathbf{D}_{n-1}\varphi(z); \mathbf{D}_{n-1}^2\varphi(z)) = 0.$$

As previously noted, the map φ on the left hand side of this equation depends on the sequence w^ν as well as the subsequence ν_k , but the right hand side is independent of these choices. Using Weierstrass theorem on the uniform convergence of derivatives, we conclude that for each sequence $\{w^\nu\}$ in U_n converging to a point in $\mathbf{b}U_n$, we have

$$\lim_{\nu \rightarrow \infty} H_j(\mathbf{D}_{n-1}f(\cdot, w^\nu); \mathbf{D}_{n-1}^2f(\cdot, w^\nu)) = 0,$$

where $\mathbf{D}_{n-1}f$ and \mathbf{D}_{n-1}^2f denote the collection of the first and second partial derivatives of f with respect to the coordinates. Consequently, we have by the maximum principle that for $(z, w) \in G_{n-1} \times U_n = G$

$$H_j(\mathbf{D}_{n-1}f(z, w); \mathbf{D}_{n-1}^2f(z, w)) = 0,$$

i.e., on G we have

$$H_j(\mathbf{D}_{n-1}f; \mathbf{D}_{n-1}^2f) \equiv 0.$$

Recalling that this equation is a restatement of (6.7), we conclude therefore that

$$\Delta_{\mu-1}(\mathbf{D}_{n-1}f) \frac{\partial}{\partial z_j} \Delta_\mu(\mathbf{D}_{n-1}f) - \Delta_\mu(\mathbf{D}_{n-1}f) \frac{\partial}{\partial z_j} \Delta_{\mu-1}(\mathbf{D}_{n-1}f) = 0,$$

which implies that

$$\frac{\partial}{\partial z_j} \left(\frac{\Delta_{\mu-1}(\mathbf{D}_{n-1}f)}{\Delta_\mu(\mathbf{D}_{n-1}f)} \right) \equiv 0 \text{ for } j \in I_{n-1},$$

holds outside the proper analytic subset $A = \{\Delta_\mu(D_{n-1}f) = 0\}$ of G . Therefore, there is a function g_n defined on $G \setminus A$, and depending only on z_n such that on $G \setminus A$, we have

$$g_n = -\frac{\Delta_{\mu-1}(D_{n-1}f)}{\Delta_\mu(D_{n-1}f)}.$$

Now let again w^ν be a sequence in U_n converging to a point of $\mathbf{b}U_n$, and let φ and c be the corresponding quantities as in (6.2). Taking setting $z_n = w^\nu$ in the above equation, using relation (6.6) and taking the limit we see that $\lim_{\nu \rightarrow \infty} g_n(w^\nu) = c$. Again, in the function on G given by $r(z) = q(f(z); g_n(z_n))$, setting $z_n = w^\nu$ and letting $\nu \rightarrow \infty$ shows that $r(z, w^\nu) \rightarrow 0$ as $\nu \rightarrow \infty$. It follows that $r(z, z_n) \rightarrow 0$ as $z_n \rightarrow \mathbf{b}U_n$. Consequently, by the maximum principle $r \equiv 0$ on $G \setminus A$, i.e., we have $q(f(z); g_n(z_n)) \equiv 0$ on $G \setminus A$. Since V is bounded, it follows that the roots of the equation $q(f(z); t) = 0$ are bounded independently of $z \in G$, so that g is a bounded holomorphic function, and therefore extends holomorphically to G . Consequently we have the relation $q(f(z); g_n(z_n)) \equiv 0$ on the whole of G .

If $\mu = 1$, we take $k = 2$ in (6.5). One can repeat the arguments to conclude again that there is a holomorphic g_n on U_n such that $q(f(z); g_n(z_n)) \equiv 0$ on G .

Repeating the argument for each factor U_j in the product $G = U_1 \times \cdots \times U_n$, we conclude that there is a function $g_j : U_j \rightarrow V$ for each $j \in I_n$ such that $q(f(z); g_j(z_j)) = 0$ for z in G . It follows now that $\pi(g_1(z_1), \dots, g_n(z_n)) = 0$. \square

We can now complete the proofs of Corollary 1.3 and Corollary 1.5:

Proof of Corollary 1.3. Since Φ and π are both proper holomorphic, $\Phi \circ \pi$ is proper holomorphic map from U^n to $\Sigma^n V$. From Theorem 1.2, we conclude that there exist proper holomorphic maps $\varphi_j : U \rightarrow V$, $j = 1, \dots, n$, such that

$$\Phi \circ \pi = \pi \circ (\varphi_1 \times \cdots \times \varphi_n).$$

We have $\Phi \circ \pi(z_1, \dots, z_n) = \pi(\varphi_1(z_1), \dots, \varphi_n(z_n))$, where $z = (z_1, \dots, z_n) \in U^n$. Hence, for each $\sigma \in S_n$ and $z \in U^n$, we get, using the symmetry of the map π that

$$\pi(\varphi_1(z_1), \dots, \varphi_n(z_n)) = \Phi \circ \pi(z) = \Phi \circ \pi(\sigma z) = \pi(\varphi_1(z_{\sigma(1)}), \dots, \varphi_n(z_{\sigma(n)})).$$

This is possible only when there is a $\varphi : U \rightarrow V$ such that

$$\varphi_1 = \varphi_2 = \cdots = \varphi_n = \varphi.$$

Hence, $\Phi = \Sigma^n \varphi$. \square

Proof of Corollary 1.5. Since $\Phi : \Sigma^n U \rightarrow \Sigma^n V$ is a proper holomorphic map, by Corollary 1.3, there is a proper holomorphic map $\varphi : U \rightarrow V$ such that $\Phi = \Sigma^n \varphi$. On the other hand proper holomorphic maps between domains with \mathcal{C}^∞ -boundaries extend to \mathcal{C}^∞ maps of the closures. This can be seen either from the general result in several variables (see [21, 7]), or in a more elementary way, by noting that the proper map being of finite degree, the branching locus is a finite set, and near the boundary the proper map is actually conformal, so that the classical theory of boundary regularity in one variable may be used. The result now follows from Theorem 1.4. \square

REFERENCES

- [1] Agler, J., Young, N. J.: A commutant lifting theorem for a domain in \mathbb{C}^2 and spectral interpolation. *J. Funct. Anal.* **161**(2), 452–477 (1999)
- [2] Agler, J., Young, N. J.: A Schwarz lemma for the symmetrized bidisc. *Bull. London Math. Soc.* **33**(2), 175–186 (2001)
- [3] Agler, J., Young, N. J.: The hyperbolic geometry of the symmetrized bidisc. *J. Geom. Anal.* **14**(3), 375–403 (2004)
- [4] Agler, J., Young, N. J.: A model theory for Γ -contractions. *J. Operator Theory* **49**(1), 45–60 (2003)
- [5] Arbarello, E., Cornalba, M., Griffiths, P. A., Harris, J.: *Geometry of algebraic curves. Vol. I.* Grundlehren der Mathematischen Wissenschaften, **267** Springer-Verlag, New York, 1985
- [6] Bedford, E., Bell, S.: Boundary continuity of proper holomorphic correspondences. *Séminaire d'analyse P. Lelong-P. Dolbeault-H. Skoda, années 1983/1984*, 47–64, *Lecture Notes in Math.*, **1198**, Springer, Berlin (1986)
- [7] Bell, S. R., Catlin, D.: Boundary regularity of proper holomorphic mappings. *Duke Math. J.* **49**(2), 385–396 (1982)
- [8] Bierstone, E., Milman, P. D.: Semianalytic and subanalytic sets. *Inst. Hautes Études Sci. Publ. Math.* **67**, 542 (1988)

- [9] Bochnak, J., Coste, M., Roy, M.-F.: Real algebraic geometry. Translated from the 1987 French original. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, 36. Springer-Verlag, Berlin (1998)
- [10] Cartan, H.: Quotient d'un espace analytique par un groupe d'automorphismes. In *Algebraic geometry and topology*. pp. 90–102. Princeton University Press, Princeton, N. J. (1957)
- [11] Cartan, H.: Quotients of complex analytic spaces. *Contributions to function theory (Internat. Colloq. Function Theory, Bombay, 1960)* pp. 1–15 Tata Institute of Fundamental Research, Bombay
- [12] Chakrabarti, D., Shaw, M.-C.: The Cauchy-Riemann equations on product domains. *Math. Ann.* **349**(4), 977–998 (2011)
- [13] Chakrabarti, D., Shaw, M.-C.: Sobolev regularity of the $\bar{\partial}$ -equation on the Hartogs triangle. *Math. Ann.* **356**(1), 241–258 (2013)
- [14] Chaumat, J., Chollet, A.-M.: Régularité höldérienne de l'opérateur $\bar{\partial}$ sur le triangle de Hartogs. *Ann. Inst. Fourier (Grenoble)* **41**(4), 867–882 (1991)
- [15] Chaumat, J., Chollet, A.-M.: Noyaux pour résoudre l'équation $\bar{\partial}$ dans des classes ultradifférentiables sur des compacts irréguliers de \mathbb{C}^n . *Several complex variables (Stockholm, 1987/1988)*, 205–226, *Math. Notes*, 38, Princeton Univ. Press, Princeton, NJ (1993)
- [16] Chen, S.-C., Shaw, M.-C. *Partial differential equations in several complex variables*. AMS/IP Studies in Advanced Mathematics, 19. American Mathematical Society, Providence, RI; International Press, Boston, MA (2001)
- [17] Conway, J. B.: *Functions of one Complex Variable II*. Graduate Texts in Mathematics, 159, Springer-Verlag, New York (1995)
- [18] Costara, C.: The symmetrized bidisc and Lempert's theorem. *Bull. London Math. Soc.* **36**(5), 656–662 (2004)
- [19] Costara, C.: The 2×2 spectral Nevanlinna-Pick problem. *J. London Math. Soc.*(2) **71**(3), 684–702 (2005)
- [20] Dufresnoy, A.: Sur l'opérateur d'' et les fonctions différentiables au sens de Whitney. *Ann. Inst. Fourier (Grenoble)* **29**(1), xvi, 229–238 (1979)
- [21] Diederich, K., Fornæss, J. E.: Boundary regularity of proper holomorphic mappings. *Invent. Math.* **67**(3), 363–384 (1982)
- [22] Edigarian, A.: Proper holomorphic self-mappings of the symmetrized bidisc. *Ann. Polon. Math.* **84**(2), 181–184 (2004)
- [23] Edigarian, A., Zwonek, W.: Geometry of the symmetrized polydisc. *Arch. Math. (Basel)* **84**(4), 364–374 (2005)
- [24] Ehsani, D.: Solution of the $\bar{\partial}$ -Neumann problem on a bi-disc. *Math. Res. Lett.* **10**(4), 523–533 (2003)
- [25] Greene, R. E., Krantz, S. G.: *Function theory of one complex variable*. Third edition. Graduate Studies in Mathematics, 40. American Mathematical Society, Providence, RI (2002)
- [26] Folland, G. B., Kohn, J. J.: *The Neumann problem for the Cauchy-Riemann complex*. Annals of Mathematics Studies, No. 75. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo (1972)
- [27] Gunning, R. C., Rossi, H.: *Analytic functions of several complex variables*. Reprint of the 1965 original. AMS Chelsea Publishing, Providence, RI (2009)
- [28] Hörmander, L.: *An introduction to complex analysis in several variables*. Third edition. North-Holland Mathematical Library, 7. North-Holland Publishing Co., Amsterdam (1990)
- [29] Kohn, J. J.: Global regularity for $\bar{\partial}$ on weakly pseudo-convex manifolds. *Trans. Amer. Math. Soc.* **181**, 273–292 (1973)
- [30] Kosiński, L.: Personal Communication, May 12, 2013.
- [31] Kurosh, A.: *Higher algebra*. Translated from the Russian by George Yankovsky. Reprint of the 1972 translation. “Mir”, Moscow (1988)
- [32] Magnus, W.: Braid groups: a survey. *Proceedings of the Second International Conference on the Theory of Groups (Australian Nat. Univ., Canberra, 1973)*, pp. 463–487. *Lecture Notes in Math.*, Vol. 372, Springer, Berlin, 1974.
- [33] Michel, J., Shaw, M.-C.: The $\bar{\partial}$ -problem on domains with piecewise smooth boundaries with applications. *Trans. Amer. Math. Soc.* **351**(11), 4365–4380 (1999)
- [34] Misra, G., Shyam Roy, S. and Zhang, G.: Reproducing kernel for a class of weighted Bergman spaces on the symmetrized polydisc. *Proc. Amer. Math. Soc.* **141**(7), 2361–2370 (2013)
- [35] Narasimhan, R.: *Several complex variables*. Reprint of the 1971 original. *Chicago Lectures in Mathematics*. University of Chicago Press, Chicago, IL (1995)
- [36] Olver, P. J.: *Classical invariant theory*. London Mathematical Society Student Texts, 44. Cambridge University Press, Cambridge, (1999)
- [37] Remmert, R., Stein, K.: Eigentliche holomorphe Abbildungen; *Math. Z.* **73**, 159–189 (1960)
- [38] Straube, E. J.: *Lectures on the L^2 -Sobolev theory of the $\bar{\partial}$ -Neumann problem*. ESI Lectures in Mathematics and Physics. European Mathematical Society (EMS), Zürich (2010).
- [39] Stout, E. L.: *Polynomial convexity*. Progress in Mathematics, 261. Birkhäuser Boston, Inc., Boston, MA (2007)
- [40] Tsuji, M.: *Potential theory in modern function theory*. Reprinting of the 1959 original. Chelsea Publishing Co., New York (1975)
- [41] Whitney, H.: *Complex analytic varieties*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. (1972)

TIFR CENTRE FOR APPLICABLE MATHEMATICS, SHARADA NAGAR, CHIKKABOMMASANDRA, BENGALURU-560065, INDIA
E-mail address: `debraj@math.tifrbng.res.in`

STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE, 8TH MILE, MYSORE ROAD, BENGALURU-560059, INDIA
E-mail address: `sushil@isibang.ac.in`